

# Mixed correlation functions in the 2-matrix model, and the Bethe ansatz

*B. Eynard*<sup>1</sup>, *N. Orantin*<sup>2</sup>

Service de Physique Théorique de Saclay,  
F-91191 Gif-sur-Yvette Cedex, France.

## Abstract:

Using loop equation technics, we compute all mixed traces correlation functions of the 2-matrix model to large  $N$  leading order. The solution turns out to be a sort of Bethe Ansatz, i.e. all correlation functions can be decomposed on products of 2-point functions. We also find that, when the correlation functions are written collectively as a matrix, the loop equation are equivalent to commutation relations.

## 1 Introduction

Formal random matrix models have been used for their interpretation as combinatorial generating functions for discretized surfaces [23, 4, 5]. The hermitean one-matrix model counts surfaces made of polygons of only one color, where as the hermitean two-matrix model counts surfaces made of polygons of two colors. In that respect, the 2-matrix model is more appropriate for the purpose of studying surfaces with non-uniform boundary conditions. At the continuum limit, the 2-matrix model gives access to “boundary operators” in conformal field theory [21].

Generating functions for surfaces with boundaries are obtained as random matrix expectation values. The expectation value of a product of  $l$  traces, is the generating function for surfaces with  $l$  boundaries, the total power of matrices in each trace being the length of the corresponding boundary. If each trace contains only one type of

---

<sup>1</sup> E-mail: [eynard@cea.fr](mailto:eynard@cea.fr)

<sup>2</sup> E-mail: [orantin@cea.fr](mailto:orantin@cea.fr)

matrix (different traces may contain different types of matrices), the expectation value is the generating function counting surfaces with uniform boundary conditions. Those non-mixed expectation values have been computed for finite  $n$  since the work of [15, 24] and refined by [1].

Mixed correlation functions have been considered as a difficult problem for a long time, and progress have been obtained only recently [2, 14]. Indeed, non-mixed expectation values can easily be written in terms of eigenvalues only (since the trace of a matrix is clearly related to its eigenvalues), where as mixed correlation functions can not ( $Tr M_1^k M_2^{k'}$  can not be written in terms of eigenvalues of  $M_1$  and  $M_2$ ).

The large  $N$  limit of the generating function of the bicolored disc (i.e. one boundary, two colors, i.e.  $\langle Tr M_1^k M_2^{k'} \rangle$ ) has been known since [19, 10, 11]. The large  $N$  limit of the generating function of the 4-colored disc (i.e. one boundary, 4 colors, i.e.  $\langle Tr M_1^k M_2^{k'} M_1^{k''} M_2^{k'''} \rangle$ ) has been known since [7]. The all order expansion of correlation functions for the 1-matrix model has been obtained by a feynman-graph representation in [8], and the generalization to non-mixed correlation functions of the 2-matrix model has been obtained in [9].

Recently, the method of integration over the unitary group of [14] has allowed to compute, for finite  $N$ , all mixed correlation functions of the 2-matrix model in terms of orthogonal polynomials.

The question of computing mixed correlation functions in the large  $N$  limit is addressed in the present article.

The answer is (not so) surprisingly related to classical results in integrable statistical models, i.e. the Bethe ansatz. It has been known for a long time that random matrix models are integrable in some sense (toda, KP, KdV, isomonodromic systems,...), but the relationship with Yang-Baxter equations and Bethe ansatz was rather indirect. Here, we see it appears in a very clear way.

outline of the article:

- section 1 is an introduction,
- in section 2, we set definitions of the model and correlation functions, and we write the relevant loop equations,
- in section 3, we introduce the Bethe Ansatz, and prove it in section 4,
- in section 5, we solve the problem under a matricial form,
- section 5 is dedicated to the special gaussian case.

## 2 The 2-matrix model, definitions and loop equations

## 2.1 Partition function

We are interested in the formal matrix integral:

$$Z := \int_{H_N^2} dM_1 dM_2 e^{-N \text{Tr} [V_1(M_1) + V_2(M_2) + M_1 M_2]} \quad (2-1)$$

where  $M_1$  and  $M_2$  are  $N \times N$  hermitean matrices, and  $dM_1$  (resp.  $dM_2$ ) is the product of Lebesgue measures of all independent real components of  $M_1$  (resp.  $M_2$ ).  $V_1(x)$  and  $V_2(y)$  are complex polynomials of degree  $d_1 + 2$  and  $d_2 + 1$ , called ‘‘potentials’’. The formal matrix integral is defined as a formal power series in the coefficients of the potentials (see [5]), computed by the usual Feynman method: consider a local extremum of  $e^{-N \text{Tr} [V_1(M_1) + V_2(M_2) + M_1 M_2]}$ , and expand the non quadratic part as a power series, and for each term of the series, perform the gaussian integration with the quadratic part. This method does not care about the convergence of the integral, or of the series, it makes sense only order by order, and it is in that sense that it can be interpreted as the generating function of discrete surfaces. All quantities in that model, have a  $1/N^2$  expansion (nearly by definition).

The extrema of  $V_1(x) + V_2(y) - xy$  are such that:

$$V_1'(x) = y \quad , \quad V_2'(y) = x \quad (2-2)$$

there are  $d_1 d_2$  solutions (indeed  $V_2'(V_1'(x)) = x$ ), which we note  $(\bar{x}_I, \bar{y}_I)$ ,  $I = 1, \dots, d_1 d_2$ . The extrema of  $\text{Tr} [V_1(M_1) + V_2(M_2) - M_1 M_2]$ , can be chosen diagonal (up to a unitary transformation), with  $\bar{x}_I$ 's and  $\bar{y}_I$ 's on the diagonal:

$$\begin{aligned} M_1 &= \text{diag}(\overbrace{\bar{x}_1, \dots, \bar{x}_1}^{n_1 \text{ times}}, \overbrace{\bar{x}_2, \dots, \bar{x}_2}^{n_2 \text{ times}}, \dots, \overbrace{\bar{x}_{d_1 d_2}, \dots, \bar{x}_{d_1 d_2}}^{n_{d_1 d_2} \text{ times}}) \\ M_2 &= \text{diag}(\overbrace{\bar{y}_1, \dots, \bar{y}_1}^{n_1 \text{ times}}, \overbrace{\bar{y}_2, \dots, \bar{y}_2}^{n_2 \text{ times}}, \dots, \overbrace{\bar{y}_{d_1 d_2}, \dots, \bar{y}_{d_1 d_2}}^{n_{d_1 d_2} \text{ times}}) \end{aligned} \quad (2-3)$$

The extremum around which we perform the expansion is thus characterized by a set of filling fractions:

$$\epsilon_I = \frac{n_I}{N} \quad , \quad \sum_{I=1}^{d_1 d_2} \epsilon_I = 1 \quad (2-4)$$

To summarize, let us say that the formal matrix integral is defined for given potentials and filling fractions.

The ‘‘one-cut’’ case, is the one where one of the filling fractions is 1, and all the others vanish. This is the case where the Feynman expansion is performed in the vicinity of only one extremum.

## 2.2 Enumeration of discrete surfaces

From those data, it is well known that formal matrix integrals are generating functions for the enumeration of discrete surfaces [5, 4, 18, 12].

For instance, in the one-cut case (expansion near an extremum  $\bar{x}, \bar{y}$ ), one has:


$$-\ln Z = \sum_G \frac{1}{\#\text{Aut}(G)} N^{\chi(G)} \left(\frac{g_2}{\delta}\right)^{n_{--}(G)} \left(\frac{\tilde{g}_2}{\delta}\right)^{n_{++}(G)} \left(\frac{-1}{\delta}\right)^{n_{+-}(G)} \prod_{i=3}^{d_1+1} g_i^{n_i(G)} \prod_{i=3}^{d_2+1} \tilde{g}_i^{\tilde{n}_i(G)} \quad (2-5)$$

where the summation is over all finite connected closed discrete surfaces made of polygons of two signs (+ and -). For such a surface (or graph)  $G$ ,  $\chi(G)$  is its Euler characteristic,  $n_i(G)$  is the number of  $i$ -gones carrying a + sign,  $\tilde{n}_i(G)$  is the number of  $i$ -gones carrying a - sign,  $n_{++}(G)$  is the number of edges separating two + polygons,  $n_{--}(G)$  is the number of edges separating two - polygons,  $n_{+-}(G)$  is the number of edges separating two polygons of different signs.  $\#\text{Aut}(G)$  is the number of automorphisms of  $G$ .

The  $g_i$ 's,  $\tilde{g}_i$ 's and  $\delta$  are defined as follows:

$$g_k := \left. \frac{\partial^k V_1(x)}{\partial x^k} \right|_{x=\bar{x}} \quad , \quad \tilde{g}_k := \left. \frac{\partial^k V_2(y)}{\partial x^k} \right|_{x=\bar{y}} \quad , \quad \delta := g_2 \tilde{g}_2 - 1 \quad (2-6)$$

Example of a discrete surface:



In the multicut case, i.e. with arbitrary filling fractions, matrix integrals can still be interpreted in terms of “foams” of surfaces, and we refer the reader to the appendix of [3] or to [13] for more details.


### 2.3 Enumeration of discrete surfaces with boundaries

Similarly, given a sequence of signs  $s_1, s_2, \dots, s_k, s_i \in \{1, 2\}$ , it is well known that the following quantity:

$$\left\langle \text{Tr} \left( \prod_{i=1}^k M_{s_i} \right) \right\rangle \tag{2-8}$$

is the generating function of discrete surfaces with one boundary of length  $k$ , whose sign of polygons on the edges are given by the sequence  $(s_1, \dots, s_k)$ .

Example of a discrete surface with boundary  $(+++++-----+++++-----)$ :



$$\langle \text{Tr} (M_1^6 M_2^5 M_1^5 M_2^6) \rangle = \sum_G$$

More generally, an expectation value of a product of  $n$  traces is the generating function for discrete surfaces with  $n$  boundaries.

In this article, we are interested only in one boundary, and to leading order in  $N$ , i.e. surfaces with the topology of a disc.

### 2.4 Master loop equation and algebraic curve

Let us define:

$$W(x) := \frac{1}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \right\rangle, \quad \tilde{W}(y) := \frac{1}{N} \left\langle \text{Tr} \frac{1}{y - M_2} \right\rangle \tag{2-10}$$

where the expectation values are formally computed as explained in the previous section, with the weight  $e^{-N \text{Tr} [V_1(M_1) + V_2(M_2) + M_1 M_2]}$ .  $W(x)$  (resp.  $\tilde{W}(y)$ ) is defined as a formal power series in its large  $x$  (resp. large  $y$ ) expansion, as well as in the expansion in the coefficients of the potentials.  $W(x)$  (resp.  $\tilde{W}(y)$ ) is a generating function for surfaces with one uniform boundary, i.e. with only sign  $+$  (resp. sign  $-$ ) polygons touching the boundary by an edge:

$$W(x) = \begin{array}{c} \text{circle} \\ + \end{array}, \quad \tilde{W}(y) = \begin{array}{c} \text{circle} \\ - \end{array} \tag{2-11}$$

We also define the following formal series:

$$Y(x) := W(x) - V_1'(x) \quad , \quad X(y) := \tilde{W}(y) - V_2'(y) \quad (2-12)$$

In addition, we define:

$$P(x, y) := \frac{1}{N} \left\langle \text{Tr} \frac{V_1'(x) - V_1'(M_1)}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle \quad (2-13)$$

$$U(x, y) := \frac{1}{N} \left\langle \text{Tr} \frac{1}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \right\rangle + x - V_2'(y) \quad (2-14)$$

$$U(x, y; x') := \left\langle \text{Tr} \frac{1}{x - M_1} \frac{V_2'(y) - V_2'(M_2)}{y - M_2} \text{Tr} \frac{1}{x' - M_1} \right\rangle - N^2 W(x') (U(x, y) - x + V_2'(y)) \quad (2-15)$$

$$E(x, y) := (V_1'(x) + y)(V_2'(y) + x) + P(x, y) - 1 \quad (2-16)$$

Notice that  $U(x, y)$  and  $U(x, y; x')$  are polynomials of  $y$  (with degree at most  $d_2 - 1$ ), and  $P(x, y)$  is a polynomial of both variables, of degree  $(d_1 - 1, d_2 - 1)$ , and  $E(x, y)$  is a polynomial of both  $x$  and  $y$ , of degree  $(d_1 + 1, d_2 + 1)$ .

It has been obtained in many articles [6, 25, 10, 11], that:

$$E(x, Y(x)) = \frac{1}{N^2} U(x, Y(x), x) \quad (2-17)$$

To large  $N$  leading order that equation reduces to an algebraic equation for  $Y(x)$ , called the ‘‘Master loop equation’’ [25]:

$$E(x, Y(x)) = 0 \quad (2-18)$$

(similarly, one also has  $E(X(y), y) = 0$ , which implies  $Y \circ X = \text{Id}$ , known as Matytsin’s equation [22]). The coefficients of  $E$ , i.e. of  $P(x, y)$  are entirely determined by the conditions  $\oint_{\mathcal{A}_i} y dx = 2i\pi\epsilon_i$ , for a choice of irreducible cycles on the algebraic curve.

The properties of that algebraic equation have been studied in many works [6, 20], here we assume that it is known.

## 2.5 Correlation functions, definitions

We define:

$$\overline{W}_k(x_1, y_1, x_2, \dots, x_k, y_k) := \frac{1}{N} \left\langle \text{Tr} \prod_{j=1}^k \frac{1}{x_j - M_1} \frac{1}{y_j - M_2} \right\rangle \quad (2-19)$$

$$\begin{aligned} & \overline{U}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\ := & \text{Pol } V_2'(y_k) \overline{W}_k(x_1, y_1, x_2, \dots, x_k, y_k) \end{aligned}$$

$$(2-20) \quad = \frac{1}{N} \left\langle \text{Tr} \frac{1}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1} \frac{V_2'(y_k) - V_2'(M_2)}{y_k - M_2} \right\rangle$$

$$(2-21) \quad \begin{aligned} & \bar{P}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\ := & \text{Pol Pol } V_1'(x_1) V_2'(y_k) \bar{W}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\ = & \frac{1}{N} \left\langle \text{Tr} \frac{V_1'(x_1) - V_1'(M_1)}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1} \frac{V_2'(y_k) - V_2'(M_2)}{y_k - M_2} \right\rangle \end{aligned}$$

$$A_k(x_1, y_1, x_2, \dots, x_k) := \frac{1}{N} \left\langle \text{Tr} \frac{1}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1} V_2'(M_2) \right\rangle \quad (2-22)$$

The functions  $\bar{W}_k$  are generating functions for discrete discs with all possible boundary conditions. One can recover any generating function of type eq. (2-8), by expanding into powers of the  $x_i$ 's and  $y_i$ 's.

For convenience, we prefer to consider the following functions:

$$W_k(x_1, y_1, x_2, \dots, x_k, y_k) := \bar{W}_k(x_1, y_1, x_2, \dots, x_k, y_k) + \delta_{k,1} \quad (2-23)$$

$$U_k(x_1, y_1, x_2, \dots, x_k, y_k) := \bar{U}_k(x_1, y_1, x_2, \dots, x_k, y_k) + \delta_{k,1}(V_2'(y_k) + x_k) \quad (2-24)$$

and for  $k > 1$ :

$$P_k(x_1, y_1, x_2, \dots, x_k, y_k) := \bar{P}_k(x_1, y_1, x_2, \dots, x_k, y_k) + W_{k-1}(x_2, \dots, x_k, y_1) \quad (2-25)$$

For the smallest values of  $k$ , those expectation values can be found in the literature, to large  $N$  leading order:

- it was found in [10, 11, 6]:

$$W_1(x, y) = \frac{E(x, y)}{(x - X(y))(y - Y(x))} \quad , \quad U_1(x, y) = \frac{E(x, y)}{(y - Y(x))} \quad (2-26)$$

- it was found in the appendix C of [7] (there is a change of sign, because the action in [7] was  $e^{-N \text{tr}(V_1(M_1) + V_2(M_2) - M_1 M_2)}$ ):

$$W_2(x_1, y_1, x_2, y_2) = \frac{W_1(x_1, y_1)W_1(x_2, y_2) - W_1(x_1, y_2)W_1(x_2, y_1)}{(x_1 - x_2)(y_1 - y_2)} \quad (2-27)$$

- For finite  $N$ , it was found in [2], and with notations explained in [2]:

$$W_1(x, y) = \det \left( 1_N + \Pi_{N-1} \frac{1}{x - Q} \frac{1}{y - P^t} \Pi_{N-1} \right) \quad (2-28)$$

- For finite  $N$ , it was found in [14] how to compute any mixed correlation function in terms of determinants involving biorthogonal polynomials, with a formula very similar to eq. (2-28).

Here, we shall find a formula for all  $W_k$ 's in the large  $N$  limit.

## 2.6 Loop equations

Loop equations are nothing but Schwinger–Dyson equations. They are obtained by writing that an integral is invariant under a change of variable, or alternatively by writing that the integral of a total derivative vanishes.

The loop equation method is well known and explained in many works [6, 25]. Here, we write for each change of variable the corresponding loop equation (we use a presentation similar to that of [6]).

In all what follows we consider  $k > 1$ .

- the change of variable:  $\delta M_2 = \frac{1}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1}$ , implies:

$$\begin{aligned}
 A_k(x_1, \dots, x_k) &= \sum_{j=1}^{k-1} \overline{W}_j(x_1, \dots, y_j) \overline{W}_{k-j}(x_k, y_j, \dots, y_{k-1}) \\
 &\quad + \frac{x_1 \overline{W}_{k-1}(x_1, y_1, \dots, y_{k-1}) - x_k \overline{W}_{k-1}(x_k, y_1, \dots, y_{k-1})}{x_1 - x_k} \\
 &= \sum_{j=1}^{k-1} W_j(x_1, \dots, y_j) W_{k-j}(x_k, y_j, \dots, y_{k-1}) \\
 &\quad - W_{k-1}(x_k, y_1, \dots, y_{k-1}) \\
 &\quad + x_k \frac{W_{k-1}(x_1, y_1, \dots, y_{k-1}) - W_{k-1}(x_k, y_1, \dots, y_{k-1})}{x_1 - x_k}
 \end{aligned} \tag{2-29}$$

- the change of variable:  $\delta M_1 = \frac{1}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1} \frac{V_2'(y_k) - V_2'(M_2)}{y_k - M_2}$ , implies:

$$\begin{aligned}
 &(Y(x_1) - y_k) \overline{U}_k(x_1, \dots, y_k) \\
 &= \sum_{j=2}^k \frac{W_{j-1}(x_1, y_1, \dots, y_{j-1}) - W_{j-1}(x_j, y_1, \dots, y_{j-1})}{x_1 - x_j} \overline{U}_{k-j+1}(x_j, y_j, \dots, x_k, y_k) \\
 &\quad + V_2'(y_k) \frac{W_{k-1}(x_1, y_1, \dots, y_{k-1}) - W_{k-1}(x_k, y_1, x_2, \dots, y_{k-1})}{x_1 - x_k} \\
 &\quad + A_k(x_1, \dots, x_k) - \overline{P}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\
 &= \sum_{j=2}^k \frac{W_{j-1}(x_1, y_1, \dots, y_{j-1}) - W_{j-1}(x_j, y_1, \dots, y_{j-1})}{x_1 - x_j} U_{k-j+1}(x_j, y_j, \dots, x_k, y_k) \\
 &\quad + \sum_{j=1}^{k-1} W_j(x_1, \dots, y_j) W_{k-j}(x_k, y_j, \dots, y_{k-1}) \\
 &\quad - P_k(x_1, y_1, x_2, \dots, x_k, y_k)
 \end{aligned} \tag{2-30}$$

- the change of variable:  $\delta M_2 = \frac{1}{x_1 - M_1} \frac{1}{y_1 - M_2} \cdots \frac{1}{x_k - M_1} \frac{1}{y_k - M_2}$ , implies:

$$\begin{aligned}
 &(X(y_k) - x_1) \overline{W}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\
 &= \sum_{j=1}^{k-1} \frac{W_{k-j}(x_{j+1}, \dots, y_k) - W_{k-j}(x_{j+1}, \dots, x_k, y_j)}{y_k - y_j} W_j(x_1, \dots, y_j) \\
 &\quad - U_k(x_1, \dots, y_k)
 \end{aligned} \tag{2-31}$$



## 2.7 Recursive determination of the correlation functions

**Theorem 2.1** *The system of equations eq. (2-30) and eq. (2-31) for all  $k$ , has a unique solution.*

In other words, if we can find some functions  $W_k, U_k, P_k$  which obey equations 2-30 and 2-31 for all  $k$ , then they are the correlation functions we are seeking.

**proof:**

$W_1, U_1$  and  $P_1$  have already been computed in the literature.

Assume that we have computed  $W_j, U_j, P_j$  for all  $j < k$ . Let us show that eqs. 2-30 and 2-31 determine uniquely  $W_k, U_k$  and  $P_k$ .

Let  $X^{(\alpha)}(y_k), \alpha = 0, \dots, d_1$  be the  $d_1 + 1$  solutions for  $x$  of  $E(x, y_k) = 0$ . I.e. for every  $\alpha = 0, \dots, d_1$  one has:

$$Y(X^{(\alpha)}(y_k)) = y_k \quad (2-32)$$

At  $x_1 = X^{(\alpha)}(y_k)$ , equation 2-30 reads:

$$\begin{aligned} & P_k(X^{(\alpha)}(y_k), y_1, x_2, \dots, x_k, y_k) \\ &= \sum_{j=2}^k \frac{W_{j-1}(X^{(\alpha)}(y_k), y_1, \dots, y_{j-1}) - W_{j-1}(x_j, y_1, x_2, \dots, y_{j-1})}{X^{(\alpha)}(y_k) - x_j} U_{k-j+1}(x_j, y_j, \dots, x_k, y_k) \\ &+ \sum_{j=1}^{k-1} W_j(X^{(\alpha)}(y_k), \dots, y_j) W_{k-j}(x_k, y_j, \dots, y_{k-1}) \end{aligned} \quad (2-33)$$

where all the quantities in the RHS are known from the recursion hypothesis. We thus know the value of  $P_k$  for  $d_1 + 1$  values of  $x_1$ . Since  $P_k$  is a polynomial in  $x_1$  of degree at most  $d_1 - 1$ , we can determine  $P_k$  by the interpolation formula:

$$\begin{aligned} & (x_1 - X(y_k)) \frac{P_k(x_1, \dots, y_k)}{E(x_1, y_k)} \\ &= \sum_{\alpha=1}^{d_2} \frac{(X^{(\alpha)}(y_k) - X(y_k)) P_k(X^{(\alpha)}(y_k), \dots, y_k)}{(x_1 - X^{(\alpha)}(y_k)) E_x(X^{(\alpha)}(y_k), y_k)} \end{aligned} \quad (2-34)$$

where  $X_k = X(y_k)$  denotes  $X^{(0)}(y_k)$ . Once  $P_k$  is known, equation 2-30 allows to compute  $U_k$ , and eq. 2-31 allows to compute  $W_k$ .  $\square$

## 3 A ‘‘Bethe ansatz’’ for correlation functions

Thus, the loop equations determine  $W_k$  uniquely, i.e., if we can find  $W_k, U_k = \text{Pol } V_2'(y_k)W_k$  and  $P_k = W_{k-1} + \text{Pol } V_1'(x_1)U_k$  which satisfy eqs. 2-31, 2-30, it means that we have the right solution. We can thus make an ansatz for  $W_k$ , and check that it satisfies the loop equations above.

Our ansatz is similar to the Bethe Ansatz [17]:

$$W_k(x_1, y_1, \dots, x_k, y_k) = \sum_{\sigma \in \Sigma_k} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \quad (3-1)$$

where the coefficients  $C_\sigma$  are rational fractions of the  $x_i$ 's and  $y_i$ 's, with at most simple poles at coinciding points, and independent of the potentials. We call eq. (3-1) a ‘‘Bethe ansatz’’, because it is very similar to the initial solution found by Bethe for the 1-dimensional spin chain, and then for the  $\delta$ -interacting bosons.

If we assume that eq. (3-1) satisfies eq. (2-31), we can in particular take the residue of eq. (2-31) at  $y_k \rightarrow Y(x_l)$  for some  $l$ . That implies the following relationship among the coefficients  $C_\sigma^{(k)}$ 's:

$$(3-2) \quad \begin{aligned} & (x_{\sigma^{-1}(k)} - x_1) C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y_k) \\ &= \sum_{j=1}^{k-1} \sum_{\tau \in \Sigma(1, \dots, j)} \sum_{\rho \in \Sigma(j+1, \dots, k)} \delta_{\sigma, \tau \rho} \frac{C_\tau^{(j)}(x_1, \dots, y_j) C_\rho^{(k-j)}(x_{j+1}, \dots, y_k)}{y_k - y_j} \end{aligned}$$

Beside, since  $W_k$  is the expectation value of a trace, the  $C_\sigma^{(k)}$ 's must be cyclically invariant:

$$(3-3) \quad C_\sigma^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = C_\sigma^{(k)}(x_2, y_2, \dots, x_k, y_k, x_1, y_1)$$

and, since  $W_k$  should have no poles at coinciding points  $y_k = y_j$  one should have:

$$(3-4) \quad \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) dy'_k = \text{Res}_{y'_k \rightarrow y_j} C_{(k,j) \circ \sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) dy'_k$$

With  $C^{(1)} = 1$ , it is clear that the set of equations eq. (3-2), eq. (3-3), eq. (3-4), have at most a unique solution. We prove in the next section that the solution exists, and thus, eq. (3-2), eq. (3-3), eq. (3-4) determine  $C_\sigma^{(k)}$  uniquely. The  $C_\sigma^{(k)}$ 's are explicitly computed in section 4.

Then in section section 4.7, we prove that:

**Theorem 3.1** *If the  $C_\sigma^{(k)}$ 's are rational functions defined by eq. (3-2), eq. (3-3), eq. (3-4), then the functions  $\hat{W}_k$ 's defined by the RHS of eq. (3-1), the functions  $\hat{U}_k(x_1, \dots, y_k) := \text{Pol}(V'_2(y_k))\hat{W}_k(x_1, \dots, y_k)$ , and the functions  $\hat{P}_k(x_1, \dots, y_k) := \text{Pol}(V'_1(x_1))\hat{U}_k(x_1, \dots, y_k) + \hat{W}_{k-1}(x_2, \dots, x_k, y_1)$ , satisfy eq. (2-31) and eq. (2-30).*

As a corollary, using theorem 2.1, we have:

**Theorem 3.2**

$$W_k(x_1, y_1, \dots, x_k, y_k) = \sum_{\sigma \in \Sigma_k} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \tag{3-5}$$

The derivation of theorem 3.1 is quite technical, and is presented in section 4.7.

## 4 Amplitudes of permutations

In this section, we compute the amplitudes  $C_\sigma^{(k)}$  explicitly.

eq. (3-2), eq. (3-3), eq. (3-4) and initial condition  $C^{(1)}(x_1, y_1) = 1$  clearly define at most a unique function  $C_\sigma^{(k)}(x_1, \dots, y_k)$ . In this section, we build the solution explicitly, and then, we prove that the function we have constructed indeed satisfies eq. (3-2), eq. (3-3), eq. (3-4).

It is easy to see that eq. (3-2) implies that  $C_\sigma^{(k)}$  vanishes for non planar permutations, and, for planar permutations,  $C_\sigma^{(k)}$  is the product of  $C_{\text{Id}}^{(k)}$  corresponding to faces. We are thus led to introduce the following definitions:

### 4.1 Some definitions: planar permutations

Let  $S$  be the shift permutation:

$$S := \text{shift} = (1, 2, \dots, k-1, k) \quad , \quad \text{i.e.} \quad S(i) = i+1 \tag{4-1}$$

**Definition 4.1** *A permutation  $\sigma \in \Sigma_k$  is called planar if*

$$n_{\text{cycles}}(\sigma) + n_{\text{cycles}}(S\sigma) = k + 1 \tag{4-2}$$


where  $n_{\text{cycles}}(\sigma)$  is the number of irreducible cycles of the permutation  $\sigma$ .

Let  $\bar{\Sigma}_k \subset \Sigma_k$  be the set of planar permutations of rank  $k$ .

eq. (4-2) is equivalent to saying that if one draws the points  $x_1, y_1, x_2, y_2, \dots, x_k, y_k$  on a circle, and draws a line between each pair  $(x_j, y_{\sigma(j)})$ , the lines don't intersect. The cycles of  $\sigma$  and the cycles of  $S\sigma$  correspond to the faces (i.e. the connected components) of that partition of the disc.

Each planar permutation can also be represented as an arch system, and thus, the number of possible planar permutations is related to Catalan number  $\text{Cat}(k)$ :

$$\text{Card}(\bar{\Sigma}_k) = \text{Cat}(k) = \frac{2k!}{k!(k+1)!} \tag{4-3}$$



Example of a planar permutation and its faces.

## 4.2 Face amplitudes

**Definition 4.2** For any  $k \geq 1$ , we define a rational function of  $x_1, \dots, y_k$ :

$$F^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k) \tag{4-5}$$

by the recursion formula:

$$\begin{aligned}
 F^{(1)}(x_1, y_1) &:= 1 \\
 F^{(k)}(x_1, y_1, \dots, x_k, y_k) &:= \sum_{j=1}^{k-1} \frac{F^{(j)}(x_1, y_1, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_k)}{(x_k - x_1)(y_k - y_j)}
 \end{aligned}
 \tag{4-6}$$

**Lemma 4.1**  $F^{(k)}$  has cyclic invariance, i.e.

$$F^{(k)}(x_2, y_2, \dots, x_k, y_k, x_1, y_1) = F^{(k)}(x_1, y_1, \dots, x_k, y_k) \tag{4-7}$$

**proof:**

We prove it by recursion. It is clearly true for  $k = 1$  and  $k = 2$  since  $F^{(2)}(x_1, y_1, x_2, y_2) = \frac{1}{(x_2 - x_1)(y_2 - y_1)}$ . For  $k \geq 3$ , assume that it is true for all  $F^{(j)}$  with  $j < k$ . One has:

$$\begin{aligned}
 & F^{(k)}(x_2, y_2, \dots, x_k, y_k, x_1, y_1) \\
 = & \sum_{j=2}^k \frac{F^{(j-1)}(x_2, \dots, x_j, y_j) F^{(k+1-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_k, x_1, y_1)}{(x_1 - x_2)(y_1 - y_j)} \\
 = & \sum_{j=2}^k \frac{F^{(j-1)}(x_2, \dots, x_j, y_j) F^{(k+1-j)}(x_1, y_1, x_{j+1}, y_{j+1}, \dots, x_k, y_k)}{(x_1 - x_2)(y_1 - y_j)}
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{j=2}^k \frac{F^{(j-1)}(x_2, \dots, x_j, y_j)}{(x_1 - x_2)(y_1 - y_j)} \\
&\quad \sum_{l=j+1}^{k-1} \frac{F^{(l+1-j)}(x_1, y_1, x_{j+1}, y_{j+1}, \dots, x_l, y_l) F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y_k)}{(x_k - x_1)(y_k - y_l)} \\
&+ \sum_{j=2}^k \frac{F^{(j-1)}(x_2, \dots, x_j, y_j)}{(x_1 - x_2)(y_1 - y_j)} \frac{F^{(1)}(x_1, y_1) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_k)}{(x_k - x_1)(y_k - y_1)} \\
&= \sum_{l=3}^{k-1} \frac{F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y_k)}{(x_k - x_1)(y_k - y_l)} \\
&\quad \sum_{j=2}^l \frac{F^{(j-1)}(x_2, \dots, x_j, y_j) F^{(l+1-j)}(x_{j+1}, y_{j+1}, \dots, x_l, y_l, x_1, y_1)}{(x_1 - x_2)(y_1 - y_j)} \\
&= \sum_{l=3}^{k-1} \frac{F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y_k) F^{(l)}(x_2, \dots, x_l, y_l, x_1, y_1)}{(x_k - x_1)(y_k - y_l)} \\
&= \sum_{l=3}^{k-1} \frac{F^{(l)}(x_1, y_1, x_2, \dots, x_l, y_l) F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y_k)}{(x_k - x_1)(y_k - y_l)} \\
&= F^{(k)}(x_1, y_1, \dots, x_k, y_k) \\
(4-8)
\end{aligned}$$

□

**Lemma 4.2** For  $k \geq 2$ ,  $F^{(k)}$  has simple poles in  $y_k$ :

$$F^{(k)}(x_1, y_1, \dots, x_k, y_k) = \sum_{l=1}^{k-1} \frac{1}{y_k - y_l} \operatorname{Res}_{y'_k \rightarrow y_l} F^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \quad (4-9)$$

**proof:**

It is clearly true for  $k = 2$ . We prove it by induction on  $k$ . Assume that it is true up to  $k - 1$ . Using the recursion hypothesis, it is clear that each term in the RHS of eq. (4-6) has at most a simple pole at  $y_k = y_l$ , and thus the recursion hypothesis is true for  $k$ . □

### 4.3 The amplitudes $C_\sigma$

**Definition 4.3** Then, for any  $k \geq 1$ , and for any permutation  $\sigma \in \Sigma_k$ , we define  $C_\sigma^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k)$  a rational function of  $x_1, \dots, y_k$ , by:

- $C_\sigma^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k) := 0$  if  $\sigma$  is not planar, and
- if  $\sigma$  is planar, we decompose  $\sigma$  and  $S\sigma$  into their product of cycles:

$$\sigma = \sigma_1 \sigma_2 \dots \sigma_l \quad , \quad S\sigma = \tilde{\sigma}_1 \tilde{\sigma}_2 \dots \tilde{\sigma}_l \quad (4-10)$$

such that:

$$\sigma_j = (i_{j,1}, i_{j,2}, \dots, i_{j,l_j}) \quad , \quad \sigma(i_{j,m}) = i_{j,m+1} \quad (4-11)$$

$$\tilde{\sigma}_j = (\tilde{i}_{j,1}, \tilde{i}_{j,2}, \dots, \tilde{i}_{j,\tilde{l}_j}) \quad , \quad \sigma(\tilde{i}_{j,m}) = \tilde{i}_{j,m+1} - 1 \quad (4-12)$$

$$C_\sigma^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k) \quad := \quad \prod_{j=1}^l F^{(l_j)}(x_{i_{j,1}}, y_{i_{j,2}}, x_{i_{j,2}}, y_{i_{j,3}}, \dots, x_{i_{j,l_j}}, y_{i_{j,1}}) \\ \prod_{j=1}^{\tilde{l}} F^{(\tilde{l}_j)}(x_{\tilde{i}_{j,1}}, y_{\tilde{i}_{j,2}-1}, x_{\tilde{i}_{j,2}}, \dots, y_{\tilde{i}_{j,\tilde{l}_j}-1}, x_{\tilde{i}_{j,\tilde{l}_j}}, y_{\tilde{i}_{j,1}-1}) \quad (4-13)$$

i.e.  $C_\sigma^{(k)}$  is the product of  $F$ 's of each connected component of the disc partitioned by  $\sigma$ .

## 4.4 Examples

### Example

In particular with  $\sigma = Id$ , we have:

$$C_{Id}^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k) = F^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \quad (4-14)$$

and with  $\sigma = S^{-1}$ :


$$C_{S^{-1}}^{(k)}(x_1, y_1, x_2, \dots, x_k, y_k) = F^{(k)}(x_k, y_{k-1}, x_{k-1}, \dots, y_1, x_1, y_k) \quad (4-15)$$

### An example with $k = 12$

Let us consider an example of  $\sigma \in \Sigma_{12}$  defined as follow:  $\sigma(1) = 3, \sigma(2) = 1, \sigma(3) = 2, \sigma(4) = 7, \sigma(5) = 6, \sigma(6) = 5, \sigma(7) = 4, \sigma(8) = 8, \sigma(9) = 11, \sigma(10) = 10, \sigma(11) = 12$  and  $\sigma(12) = 9$ . i.e.

$$\sigma = (1, 3, 2)(4, 7)(5, 6)(8)(9, 12, 11)(10) \\ S\sigma = (1, 4, 8, 9)(2)(3)(5, 7)(6)(10, 11)(12) \quad (4-16)$$

The corresponding arch system is:



where dark faces (resp. white faces) correspond to the cycles of  $\sigma$  (resp.  $S\sigma$ ). For that permutation  $C_\sigma^{(12)}$  is worth:

$$F^{(3)}(x_1, y_3, x_3, y_2, x_2, y_1)F^{(2)}(x_5, y_6, x_6, y_5)F^{(3)}(x_9, y_{12}, x_{12}, y_{11}, x_{11}, y_9) \\ \times F^{(2)}(x_4, y_7, x_7, y_4)F^{(4)}(x_1, y_3, x_4, y_7, x_8, y_8, x_9, y_{12})F^{(2)}(x_5, y_6, x_7, y_4)F^{(2)}(x_{10}, y_{10}, x_{11}, y_9) \quad (4-18)$$

**Example  $k \leq 3$**

$$C^{(1)} = 1 \quad (4-19)$$

$$C_{Id}^{(2)} = \frac{1}{(x_2 - x_1)(y_2 - y_1)} \quad (4-20)$$

$$C_{(12)}^{(2)} = \frac{1}{(x_2 - x_1)(y_1 - y_2)} \quad (4-21)$$

$$C_{Id}^{(3)} = \frac{1}{(x_1 - x_3)} \left( \frac{1}{y_3 - y_1} \frac{1}{(x_2 - x_3)(y_3 - y_2)} + \frac{1}{y_3 - y_2} \frac{1}{(x_1 - x_2)(y_2 - y_1)} \right) \quad (4-22)$$

$$C_{(1)(23)}^{(3)} = \frac{1}{(x_1 - x_2)} \frac{1}{y_3 - y_1} \frac{1}{(x_2 - x_3)(y_2 - y_3)} \quad (4-23)$$

$$C_{(12)(3)}^{(3)} = \frac{1}{(x_1 - x_3)} \frac{1}{y_3 - y_2} \frac{1}{(x_1 - x_2)(y_1 - y_2)} \quad (4-24)$$

$$C_{(123)}^{(3)} = \frac{1}{(x_1 - x_2)(x_2 - x_3)(y_1 - y_2)} \frac{1}{y_3 - y_1} + \frac{1}{(x_1 - x_3)(x_1 - x_2)(y_2 - y_1)} \frac{1}{y_3 - y_2} \quad (4-25)$$

$$C_{(13)(2)}^{(3)} = -\frac{1}{y_3 - y_1} \frac{1}{(x_1 - x_3)(x_2 - x_3)(y_1 - y_2)} \quad (4-26)$$

$$C_{(132)}^{(3)} = 0 \quad (4-27)$$

### Example: Rainbows

The rainbow is the permutation

$$\sigma(j) = k + 1 - j \quad (4-28)$$

if  $k$  is even:

$$C_{\sigma}^{(k)} = \frac{1}{\prod_{i=1}^{k/2} (x_{k+1-i} - x_i)(y_i - y_{k+1-i}) \prod_{i=1}^{k/2-1} (x_{k+1-i} - x_{i+1})(y_i - y_{k-i})} \quad (4-29)$$

if  $k$  is odd:

$$C_{\sigma}^{(k)} = \frac{1}{\prod_{i=1}^{(k-1)/2} (x_{k+1-i} - x_i)(x_{k+1-i} - x_{i+1})(y_i - y_{k+1-i})(y_i - y_{k-i})} \quad (4-30)$$

## 4.5 Properties of $C_{\sigma}$

**Lemma 4.3** *The  $C_{\sigma}^{(k)}$ 's are cyclically invariant:*

$$C_{\sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = C_{\sigma}^{(k)}(x_2, y_2, \dots, x_k, y_k, x_1, y_1) \quad (4-31)$$

**proof:**

It follows from Lemma 4.1.  $\square$

**Lemma 4.4** *The  $C_{\sigma}^{(k)}$ 's have at most simple poles in all their variables, and are such that:*

$$C_{\sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = \sum_{j=1}^{k-1} \frac{1}{y_k - y_j} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) dy'_k \quad (4-32)$$

**proof:**

If  $\sigma$  is planar, the pair  $(y_k, y_j)$  can appear in at most one factor of eq. (4-13), and the results follows from lemma 4.2.  $\square$

**Theorem 4.1** *The  $C_{\sigma}^{(k)}$ 's, with  $\sigma(1) \neq k$ , satisfy the recursion formula eq. (3-2):*

$$C_{\sigma}^{(k)} = \frac{1}{x_{\sigma^{-1}(k)} - x_1} \sum_{j=1}^{k-1} \sum_{\tau \in \Sigma(1, \dots, j)} \sum_{\rho \in \Sigma(j+1, \dots, k)} \delta_{\sigma, \tau \rho} \frac{1}{y_k - y_j} C_{\tau}^{(j)} C_{\rho}^{(k-j)} \quad (4-33)$$



**proof:**

Let  $\pi$ , be the cycle of length  $l + 1 \geq 2$  of  $S\sigma$  which contains  $x_1$  and  $y_k$ :

$$\pi = (x_1 \rightarrow y_{i_1-1} \rightarrow x_{i_1} \rightarrow y_{i_2-1} \rightarrow \dots \rightarrow y_{i_l-1} \rightarrow x_{i_l} \rightarrow y_k \rightarrow x_1) \quad (4-34)$$

$$i_{j+1} = \sigma(i_j) + 1 \quad , \quad i_0 := 1 \quad (4-35)$$

Planarity implies that:

$$i_0 < i_1 < i_2 < \dots < i_l \quad (4-36)$$

Since  $\sigma$  is planar, there exists a unique way of factorizing  $\sigma$  as:

$$\sigma = \prod_{j=0}^l \sigma_j \quad , \quad \sigma_j \in \Sigma(i_j, \dots, i_{j+1} - 1) \quad (4-37)$$

From the definition of  $C_\sigma^{(k)}$  we have:

$$C_\sigma^{(k)} = F^{(l+1)}(x_1, y_{i_1-1}, x_{i_1}, y_{i_2-1}, \dots, y_{i_l-1}, x_{i_l}, y_k) \prod_j C_{\sigma_j}^{(i_{j+1}-i_j)}(x_{i_j}, \dots, y_{i_{j+1}-1}) \quad (4-38)$$

and using 4-6, we have:

$$\begin{aligned} C_\sigma^{(k)} &= F^{(l+1)}(x_1, y_{i_1-1}, x_{i_1}, y_{i_2-1}, \dots, y_{i_l-1}, x_{i_l}, y_k) \prod_j C_{\sigma_j}^{(i_{j+1}-i_j)}(x_{i_j}, \dots, y_{i_{j+1}-1}) \\ &= \sum_{m=1}^l \frac{F^{(m)}(x_1, y_{i_1-1}, x_{i_1}, \dots, x_{i_m-1}, y_{i_m-1}) F^{(l+1-m)}(x_{i_m}, y_{i_m+1-1}, x_{i_m+1}, \dots, x_{i_l}, y_k)}{(x_{i_l}-x_1)(y_k-y_{i_m-1})} \\ &\quad \prod_j C_{\sigma_j}^{(i_{j+1}-i_j)}(x_{i_j}, \dots, y_{i_{j+1}-1}) \end{aligned} \quad (4-39)$$

notice that  $i_l = \sigma^{-1}(k)$ , and note:

$$\tau_m := \prod_{j=1}^{m-1} \sigma_j \quad , \quad \rho_m := \prod_{j=m}^l \sigma_j \quad (4-40)$$

We have:

$$\tau_m \in \Sigma(1, \dots, i_m - 1) \quad , \quad \rho_m \in \Sigma(i_m, \dots, k) \quad (4-41)$$

4-38 gives:

$$C_\sigma^{(k)} = \frac{1}{x_{\sigma^{-1}(k)} - x_1} \sum_{m=1}^l \frac{1}{y_k - y_{i_m-1}} C_{\tau_m}^{(i_m)} C_{\rho_m}^{(k-i_m)} \quad (4-42)$$

It is clear, from the planarity condition that if there exists some  $j$  and  $\tau$  and  $\rho$ , such that:

$$\sigma = \tau\rho \quad , \quad \tau \in \Sigma(1, \dots, j) \quad , \quad \rho \in \Sigma(j + 1, \dots, k) \quad (4-43)$$

then, one must have  $j = i_m$ ,  $\tau = \tau_m$  and  $\rho = \rho_m$  for some  $m$ .  $\square$

**Lemma 4.5** For any transposition  $(k, j)$  (with  $k \neq j$ ), we have:

$$\operatorname{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) dy'_k = - \operatorname{Res}_{y'_k \rightarrow y_j} C_{(k,j) \circ \sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) dy'_k \quad (4-44)$$

**proof:**

It is trivial if  $y_j$  and  $y_k$  are not in the same face, because both sides vanish: the LHS has no pole, and the RHS is a non-planar permutation. The case where  $y_j$  and  $y_k$  belong to the same face reduces to proving the Lemma for  $\sigma = Id$ .

For  $\sigma = Id$ , we prove it by recursion on  $k$ . It clearly works for  $k = 1$  and  $k = 2$ . Assume that it works up to  $k - 1$ .

From the definition 4.3 we have:

$$\begin{aligned} & C_{(k,j)}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \\ = & F^{(j)}(x_1, y_1, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, y_{k-1}, x_k, y_j) F^{(2)}(x_j, y_k, x_k, y_j) \\ = & \frac{F^{(j)}(x_1, y_1, \dots, x_j, y_k) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, y_{k-1}, x_k, y_j)}{(x_j - x_k)(y_k - y_j)} \end{aligned} \quad (4-45)$$

and thus:

$$\operatorname{Res}_{y'_k \rightarrow y_j} C_{(k,j)}^{(k)}(x_1, y_1, \dots, x_k, y_k) = \frac{F^{(j)}(x_1, y_1, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)}{(x_j - x_k)} \quad (4-46)$$

On the LHS, we have from eq. (4-6):

$$\begin{aligned} & \operatorname{Res}_{y'_k \rightarrow y_j} F^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) \\ = & \operatorname{Res}_{y'_k \rightarrow y_j} \sum_{l=1}^{k-1} \frac{F^{(l)}(x_1, y_1, x_2, \dots, x_l, y_l) F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y'_k)}{(x_k - x_1)(y'_k - y_l)} \\ = & \frac{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)}{(x_k - x_1)} \\ & + \sum_{l=1}^{j-1} \frac{F^{(l)}(x_1, y_1, x_2, \dots, x_l, y_l)}{(x_k - x_1)(y_j - y_l)} \operatorname{Res}_{y'_k \rightarrow y_j} F^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y'_k) \end{aligned} \quad (4-47)$$

Then, from the recursion hypothesis, and from eq. (4-46) we have:

$$= \frac{\operatorname{Res}_{y'_k \rightarrow y_j} F^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k)}{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)} \frac{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)}{(x_k - x_1)}$$

$$\begin{aligned}
& - \sum_{l=1}^{j-1} \frac{F^{(l)}(x_1, y_1, x_2, \dots, x_l, y_l)}{(x_k - x_1)(y_j - y_l)} \operatorname{Res}_{y'_k \rightarrow y_j} C_{(k,j)}^{(k-l)}(x_{l+1}, y_{l+1}, \dots, x_k, y'_k) \\
= & \frac{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)}{(x_k - x_1)} \\
& - \sum_{l=1}^{j-1} \frac{F^{(l)}(x_1, y_1, x_2, \dots, x_l, y_l)}{(x_k - x_1)(y_j - y_l)} \\
& \frac{F^{(j-l)}(x_{l+1}, y_{l+1}, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, y_{k-1}, x_k, y_j)}{(x_j - x_k)}
\end{aligned}
\tag{4-48}$$

In the last line, we use again eq. (4-6), and thus:

$$\begin{aligned}
& \operatorname{Res}_{y'_k \rightarrow y_j} F^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y'_k) \\
= & \frac{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j) F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, x_k, y_j)}{(x_k - x_1)} \\
& - (x_j - x_1) \frac{F^{(j)}(x_1, y_1, x_2, \dots, x_j, y_j)}{(x_k - x_1)} \frac{F^{(k-j)}(x_{j+1}, y_{j+1}, \dots, y_{k-1}, x_k, y_j)}{(x_j - x_k)}
\end{aligned}
\tag{4-49}$$

It is then clear to see that the sum of eq. (4-46) and eq. (4-49) vanishes, and the recursion hypothesis is proven for  $k$ .  $\square$

**Lemma 4.6**

$$\forall k > 1, \quad \sum_{\sigma \in \Sigma_k} C_{\sigma}^{(k)}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = 0 \tag{4-50}$$

**proof:**

This expression is a rational function of all its variables. Consider the poles at  $y_k = y_j$ , and write  $\tau = (k, j)$ . One can split the symmetric group  $\Sigma_k$  into its two conjugacy classes wrt the subgroup generated by  $\tau$ :  $\Sigma_k = [Id] \oplus [\tau]$ . In other words:

$$\begin{aligned}
& \sum_{\sigma \in \Sigma_k} C_{\sigma}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \\
= & \sum_{\sigma \in \Sigma_k / \tau} C_{\sigma}(x_1, y_1, x_2, y_2, \dots, x_k, y_k) + C_{\tau\sigma}(x_1, y_1, x_2, y_2, \dots, x_k, y_k)
\end{aligned}
\tag{4-51}$$

From Lemma 4.5, the terms in the RHS have no pole at  $y_k = y_j$ . Similarly, using cyclicity and doing the same for the  $x$ 's, we prove the lemma.  $\square$

The Lemmas we have just proven, are sufficient to prove the main theorem 3.2. This is done in section 4.7.


### 4.6 Computation of the rational functions $F^{(k)}(x_1, y_1, \dots, x_k, y_k)$

Although the exact computation of the  $F^{(k)}$ 's is not necessary for proving theorem 3.2, we do it for completeness. In this section we give an explicit (and non-recursive) formula for the  $F^{(k)}$ 's.



**Definition 4.4** Let  $\mathcal{T}_k$  be the set of planar binary skeleton trees defined as follows: A tree  $T$  belongs to  $\mathcal{T}_k$ , and is called a  $k$ -planar tree if and only if:

- it is composed of  $k+1$  vertices, labeled by  $[1, \dots, k+1]$ ;
- it has  $k$  edges which can be either upgoing or downgoing;
- its vertices have valence 1, 2 or 3, and are of one of the following eight possibilities, in which the point  $m$  denotes the origin of the branch containing  $i$ :


– two trivalent vertices :



– four bivalent vertices :



– two monovalent vertices corresponding to the leaves of the tree:



**Remark 4.1** One can see that the root is necessarily labeled  $k+1$ , and that the first edge is necessarily upgoing, and its extremity is necessarily 1.



**Theorem 4.2** There is a bijection between  $\mathcal{T}_k$  and the set of planar permutations  $\bar{\Sigma}_k$ :

**proof:**

We build explicitly that bijection.

Consider a planar permutation  $\sigma \in \overline{\Sigma}_{k-1}$ . Planarity means that  $\sigma$  defines a partition of the disc into faces of two kinds. Let us say that faces which correspond to cycles of  $S\sigma$  are colored in white, faces which correspond to cycles of  $\sigma$  are colored in black.

Decompose  $\sigma$  and  $S\sigma$  into products of irreducible cycles:

$$\sigma = \sigma_1 \sigma_2 \dots \sigma_l \quad , \quad S\sigma = \tilde{\sigma}_1 \tilde{\sigma}_2 \dots \tilde{\sigma}_{\tilde{l}} \quad (4-57)$$

and we assume that  $\tilde{\sigma}_1$  and  $\sigma_1$  contain  $x_1$ .

Because of planarity, we can define a distance of faces (i.e. cycles) to the face  $\tilde{\sigma}_1$ , as the number of edges one has to cross for going from a face  $\sigma_i$  or  $\tilde{\sigma}_i$  to  $\sigma_1$ , and call it  $D(\sigma_i)$  or  $D(\tilde{\sigma}_i)$ .

We also define the ‘‘origin’’ of a face, noted  $m(\sigma_i)$  or  $m(\tilde{\sigma}_i)$ , as follows: If the face is  $\tilde{\sigma}_1$ , we define  $m(\tilde{\sigma}_1) = k$ , otherwise, because of planarity, there is only one neighbouring face which is at smaller distance of  $\tilde{\sigma}_1$ . Because of planarity, those two faces share at most one  $x$ , and the origin is defined as the label of that  $x$ .

Thus, each face has a color, white or black, a distance  $D$ , and an origin  $m$ .

Now, to every face we associate a branch as follows:

- to a white face,  $\tilde{\sigma}_j$ , i.e. a cycle of  $S\sigma$ , noted

$$\tilde{\sigma}_j = (\tilde{i}_{j,1}, \tilde{i}_{j,2}, \dots, \tilde{i}_{j,\tilde{l}_j}) \quad , \quad \tilde{i}_{j,1} = m(\tilde{\sigma}_j) \quad , \quad \sigma(\tilde{i}_{j,n}) = \tilde{i}_{j,n+1} - 1 \quad (4-58)$$

we associate the upgoing branch  $\tilde{i}_{j,1} \rightarrow \tilde{i}_{j,2} \rightarrow \dots \rightarrow \tilde{i}_{j,\tilde{l}_j}$




(if  $\tilde{l}_j = 1$ , the sequence contains only one vertex  $\tilde{i}_{j,1} = m(\tilde{\sigma}_j)$  and no edge).

- to a black face,  $\sigma_j$ , i.e. a cycle of  $\sigma$ , noted

$$\sigma_j = (i_{j,1}, i_{j,2}, \dots, i_{j,l_j}) \quad , \quad i_{j,1} = m(\sigma_j) \quad , \quad \sigma(i_{j,n}) = i_{j,n+1} \quad (4-59)$$

we associate the downgoing branch  $i_{j,1} \rightarrow i_{j,2} \rightarrow \dots \rightarrow i_{j,l_j}$




(if  $l_j = 1$ , the sequence contains only one vertex  $i_{j,1} = m(\sigma_j)$  and no edge).

- to the first face  $\tilde{\sigma}_1$ ,

$$\tilde{\sigma}_1 = (\tilde{i}_{1,1}, \tilde{i}_{1,2}, \dots, \tilde{i}_{1,\tilde{l}_1}) \quad , \quad \tilde{i}_{1,1} = 1 \quad , \quad \sigma(\tilde{i}_{1,n}) = \tilde{i}_{1,n+1} - 1 \quad (4-60)$$

we associate the upgoing branch  $k \rightarrow \tilde{i}_{1,1} \rightarrow \tilde{i}_{1,2} \rightarrow \dots \rightarrow \tilde{i}_{1,\tilde{l}_1}$



By definition of the origin  $m$  of a face at distance  $D$ , the origin of a branch is necessarily a vertex on a branch at distance  $D - 1$ , and from planarity, it cannot be a vertex on any other branch. Thus, there is a unique way to attach all branches to their origin, and we obtain a tree, which belongs to  $\mathcal{T}_{k-1}$ .

Inverse bijection:

On the other hand, let us consider a  $k - 1$ -tree. One can build a permutation  $\sigma \in \Sigma_{k-1}$  as follows: the image of an element of  $(1, \dots, k - 1)$  is :


- its descendant along a downgoing propagator if it exists;
- the origin of the downgoing branch to which it belongs in the other cases.

Because of the form of the vertices, the upgoing branches are necessarily the cycles of  $S\sigma$ . And since the branches form a tree, it implies that two faces touch one another through zero or one edge, thus the permutation  $\sigma$  is planar.


It is easy to see that this application is the inverse of the preceding one.  $\square$

**Example:**

Let us carry out explicitly step by step this building for the permutation  $\sigma \in \Sigma_{12}$  introduced earlier. Notice that it is enrooted in  $12 + 1 = 13$ .





Considering the last non trivial cycle  $\tilde{\sigma}_6 = (10, 11)$ , one obtains finally the tree corresponding to 4-17:



(4-62)

**Corollary 4.1**  $\#\mathcal{T}_k = \text{Cat}(k)$ , where  $\text{Cat}(k)$  is the  $k$ 'th Catalan number.

**Definition 4.5** To every permutation  $\sigma \in \Sigma_{k-1}$ , we associate a weight  $f_\sigma$  computed as follows:

- To every downgoing edge  of a cycle  $\sigma_n$  of  $\sigma$ , one associates the weight  $g_{i_{n,1}, i_{n,j+1}, i_{n,j+2}}$ ;
- To every upgoing edge  of a cycle  $\tilde{\sigma}_n$  of  $S\sigma$ , one associates the weight  $g_{\tilde{i}_{n,j+1}, \tilde{i}_{n,1}, \sigma(\tilde{i}_{n,j+1})}$ .

where  $g_{i,h,j}$  is defined as follows:

$$g_{i,h,j} := \frac{1}{x_h - x_i} \frac{1}{y_h - y_j} \quad (4-63)$$

and  $f_\sigma$  is the product of all the weights of edges composing the tree associated to  $\sigma$  in  $\mathcal{T}_{k-1}$ . That is to say:

$$f_\sigma = \prod_{n=1}^l \prod_{j=2}^{l_n} g_{i_{n,1}, i_{n,j}, i_{n,j+1}} \prod_{n=2}^{\tilde{l}} \prod_{j=2}^{\tilde{l}_n} g_{\tilde{i}_{n,j}, \tilde{i}_{n,1}, \sigma(\tilde{i}_{n,j})} \prod_{j=1}^{\tilde{l}_1} g_{\tilde{i}_{1,j}, k, \sigma(\tilde{i}_{1,j})} \quad (4-64)$$

**Theorem 4.3**  $F^{(k)}(x_1, y_1, \dots, x_k, y_k)$  is obtained as the sum of the weights  $f_\sigma$ 's over all the  $k-1$ -trees:

$$F^{(k)}(x_1, y_1, \dots, x_k, y_k) = \sum_{\sigma \in \bar{\Sigma}_{k-1}} f_\sigma \quad (4-65)$$

proof:

First of all, let us interpret diagrammatically the recursion relation eq. (4-6) defining the  $F$ 's:

$$(4-66)$$

Actually, this recursion relation is nothing else but a rule for cutting a graph along the dashed line into two smaller ones. The weight of a graph is then obtained as the sum over all the possible ways of cutting it in two.

Notice that  $F^{(k)}$  is the sum of  $\text{Cat}(k - 1)$  different terms.

Let us now explicit this bijection with the graphs with  $k - 1$  arches. In order to compute one of the terms composing  $F^{(k)}$ , one has to cut it with the help of the recursion relation until one obtains only graphs with one arch. That is to say that one cuts it  $k - 1$  times along non intersecting lines (corresponding to the dashed one in the recursion relation). If one draws these cutting lines on the circle, one obtains a graph with  $k - 1$  arches dual of the original one. Thus every way of cutting a graph with  $k - 1$  arches is associated to a planar permutation  $\sigma \in \Sigma(i \dots k - 1)$ . Let us now prove that the term obtained by this cutting is equal to  $f_\sigma$ .

For the sake of simplicity, one denotes the identity graph of  $(x_j, y_j, \dots, x_k, y_k)$  by circle  $(j, j + 1, \dots, k)$ . In these conditions, the recursion relation reads:

$$(1, 2, \dots, k) = \sum_{j=1}^{k-1} g_{1,k,j}(1, \dots, j)(j + 1, \dots, k) \quad (4-67)$$

Let  $\sigma$  be a permutation of  $(1, \dots, k - 1)$ . Cut it along the line going from the boundary  $(x_1, y_k)$  to  $(y_{\sigma(1)}, x_{S\sigma(1)})$ . It results from this operation the factor  $g_{1,k,\sigma(1)}$  and the circles  $(1, \dots, \sigma(1))$  and  $(S\sigma(1), \dots, k)$ :

$$(1, \dots, k) \rightarrow^\sigma g_{1,k,\sigma(1)}(1, \dots, \sigma(1))(S\sigma(1), \dots, k) \quad (4-68)$$

Then cutting the circle  $(S\sigma(1), \dots, k)$  along  $(y_k, x_{S\sigma(1)}) \rightarrow (y_{\sigma S\sigma(1)}, x_{S\sigma S\sigma(1)})$  gives:

$$(S\sigma(1), \dots, k) \rightarrow^\sigma g_{S\sigma(1),k,\sigma S\sigma(1)}(S\sigma(1), \dots, \sigma S\sigma(1))(S\sigma S\sigma(1), \dots, k) \quad (4-69)$$

One pursues this procedure step by step by always cutting the circle containing  $k$ . Using the former notations, this reads:

$$(1, \dots, k) \rightarrow^\sigma \prod_{j=1}^{\tilde{l}_1} g_{\tilde{i}_{1,j},k,\sigma(\tilde{i}_{1,j})}(\tilde{i}_{1,j}, \dots, \sigma(\tilde{i}_{1,j})) \quad (4-70)$$



So one has computed the weight associated to the first  $S\sigma$  - cycle. The remaining circles correspond to  $\sigma$ -cycles. Let us compute their weight by considering for example  $(\tilde{i}_{1,1}, \dots, \sigma(\tilde{i}_{1,1})) = (i_{1,1}, \dots, i_{1,2})$ .

The cut along the line  $(x_{i_{1,1}}, y_{i_{1,2}}) \rightarrow (y_{i_{1,3}}, x_{S(i_{1,1})})$  gives:

$$(i_{1,1}, \dots, i_{1,2}) \rightarrow^\sigma g_{i_{1,1}, i_{1,2}, i_{1,3}}(i_{1,1}, \dots, i_{1,3})(S(i_{1,3}), \dots, i_{1,2}) \quad (4-71)$$

Keeping on cutting the circle containing  $i_{1,1}$  at every step gives:

$$(i_{1,1}, \dots, i_{1,2}) \rightarrow^\sigma \prod_{j=2}^{l_1} g_{i_{1,1}, i_{1,j}, i_{1,j+1}}(S(i_{1,j+1}), \dots, i_{1,j}) \quad (4-72)$$

One can notice that the remaining circles in the RHS correspond to cycles of  $S\sigma$  whose contribution has not been taken into account yet. One can then compute their values by following a procedure similar to the one used for the first  $S\sigma$ -cycle.


One can then recursively cut the circles so that one finally obtains only circles containing only one element. This recursion is performed by alternatively processing on  $\sigma$ -cycles and  $S\sigma$ -cycles.

Thus, one straightforwardly finds:

$$(1, \dots, k) \rightarrow^\sigma \prod_{n=1}^l \prod_{j=2}^{l_n} g_{i_{n,1}, i_{n,j}, i_{n,j+1}} \prod_{n=2}^{\tilde{l}} \prod_{j=2}^{\tilde{l}_n} g_{\tilde{i}_{n,j}, \tilde{i}_{n,1}, \sigma(\tilde{i}_{n,j})} \prod_{j=1}^{\tilde{l}_1} g_{\tilde{i}_{1,j}, k, \sigma(\tilde{i}_{1,j})} = f_\sigma \quad (4-73)$$

□


**Example:** Let us compute the weight associated to our example. Starting from the circle  $(1, \dots, 13)$ , one will proceed step by step the following cutting:



The first step consists in cutting along the  $\tilde{\sigma}_1$  cycle. The dashed lines show where one cuts the circles. Note that one do not represent the circles of unite length. The associated weight is  $g_{1,13,3} g_{4,13,7} g_{8,13,8} g_{9,13,12}$ .

The second step consists in cutting along the remaining  $\sigma$  cycles. One associates the weight  $g_{1,3,2} g_{1,2,1} \times g_{4,7,4} \times g_{9,12,11} g_{9,11,9}$  to this step.

The weights associated to the two last cuttings are  $g_{5,7,6} \times g_{10,11,10}$  and  $g_{5,6,5}$ .



(4-75)

Finally, the weight of this planar permutation is then:

$$f_{\sigma} = g_{1,13,3} g_{4,13,7} g_{8,13,8} g_{9,13,12} g_{1,3,2} g_{1,2,1} g_{4,7,4} g_{9,12,11} g_{9,11,9} g_{5,7,6} g_{10,11,10} g_{5,6,5} \quad (4-76)$$

## 4.7 Proof of the ansatz

proof of theorem 3.1:

Using eq. (4-32), one has:

$$\begin{aligned}
\hat{U}_k(x_1, \dots, y_k) &= \text{Pol}_{y_k} V_2'(y_k) \hat{W}_k(x_1, y_1, \dots, x_k, y_k) \\
&= \text{Pol}_{y_k} V_2'(y_k) \sum_{\sigma \in \Sigma_k} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \\
&= \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
&\quad \text{Pol}_{y_k} \frac{V_2'(y_k) W_1(x_{\sigma^{-1}(k)}, y_k)}{y_k - y_j} \\
&= \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
&\quad \text{Pol}_{y_k} \frac{(\hat{W}(y_k) - X(y_k)) W_1(x_{\sigma^{-1}(k)}, y_k)}{y_k - y_j} \\
&= - \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
&\quad \text{Pol}_{y_k} \frac{X(y_k) W_1(x_{\sigma^{-1}(k)}, y_k)}{y_k - y_j} \\
&= \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
&\quad \text{Pol}_{y_k} \frac{(x_{\sigma^{-1}(k)} - X(y_k)) W_1(x_{\sigma^{-1}(k)}, y_k) - (x_{\sigma^{-1}(k)} - X(y_j)) W_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j} \\
&= \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \text{Res}_{y'_k \rightarrow y_j} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y'_k) dy'_k \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
&\quad \text{Pol}_{y_k} \frac{(x_{\sigma^{-1}(k)} - X(y_k)) W_1(x_{\sigma^{-1}(k)}, y_k) - (x_{\sigma^{-1}(k)} - X(y_j)) W_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j}
\end{aligned} \tag{4-77}$$

Indeed, using eq. (2-26), one sees that the last expression is a polynomial in  $y_k$ :

$$\begin{aligned}
&\frac{(x_{\sigma^{-1}(k)} - X(y_k)) W_1(x_{\sigma^{-1}(k)}, y_k) - (x_{\sigma^{-1}(k)} - X(y_j)) W_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j} \\
&= \frac{\frac{E(x_{\sigma^{-1}(k)}, y_k)}{y_k - Y(x_{\sigma^{-1}(k)})} - \frac{E(x_{\sigma^{-1}(k)}, y_j)}{y_j - Y(x_{\sigma^{-1}(k)})}}{y_k - y_j}
\end{aligned} \tag{4-78}$$

We have to check eq. (2-31), i.e. that  $A = 0$  with:

$$\begin{aligned}
A &:= \sum_j \frac{\hat{W}_{k-j}(x_{j+1}, \dots, y_k) - \hat{W}_{k-j}(x_{j+1}, \dots, x_k, y_j)}{y_k - y_j} \hat{W}_j(x_1, \dots, y_j) \\
&\quad - \hat{U}_k(x_1, \dots, y_k) \\
&\quad + (x_1 - X(y_k)) \hat{W}_k(x_1, y_1, x_2, \dots, x_k, y_k)
\end{aligned} \tag{4-79}$$

We have:

$$\begin{aligned}
A &= \sum_{j \neq k} \sum_{\pi} \sum_{\tau} \frac{C_\tau^{(j)}(x_1, \dots, y_j) C_\pi^{(k-j)}(x_{j+1}, \dots, y_k)}{y_k - y_j} \\
&\quad \times \prod_{i=1}^j W_1(x_{\tau^{-1}(i)}, y_i) \prod_{i=j+1}^k W_1(x_{\pi^{-1}(i)}, y_i) \\
&\quad - \sum_{j \neq k} \sum_{\pi} \sum_{\tau} \frac{C_\tau^{(j)}(x_1, \dots, y_j) C_\pi^{(k-j)}(x_{j+1}, \dots, y_j)}{y_k - y_j} W_1(x_{\pi^{-1}(k)}, y_j)
\end{aligned}$$

$$\begin{aligned}
& \times \prod_{i=1}^j W_1(x_{\tau^{-1}(i)}, y_i) \prod_{i=j+1}^{k-1} W_1(x_{\pi^{-1}(i)}, y_i) \\
& - \sum_{\sigma} \sum_{j \neq k} \frac{(x_{\sigma^{-1}(k)} - X(y_k))}{y_k - y_j} W_1(x_{\sigma^{-1}(k)}, y_k) \\
& \times \left( \operatorname{Res}_{y_k \rightarrow y_j} C_{\sigma}^{(k)} \right) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& + \sum_{\sigma} \sum_{j \neq k} \frac{(x_{\sigma^{-1}(k)} - X(y_j))}{y_k - y_j} W_1(x_{\sigma^{-1}(k)}, y_j) \\
& \times \left( \operatorname{Res}_{y_k \rightarrow y_j} C_{\sigma}^{(k)} \right) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& + (x_1 - X(y_k)) \sum_{\sigma} C_{\sigma}^{(k)}(x_1, \dots, y_k) W_1(x_{\sigma^{-1}(k)}, y_k) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i)
\end{aligned}
\tag{4-80}$$

Using eq. (4-32) in the last line, adding it to the 4th line, and using eq. (4-32) again, we get:

$$\begin{aligned}
A & = \sum_{j \neq k} \sum_{\pi} \sum_{\tau} \frac{C_{\tau}^{(j)}(x_1, \dots, y_j) C_{\pi}^{(k-j)}(x_{j+1}, \dots, y_k)}{y_k - y_j} \\
& \times \prod_{i=1}^j W_1(x_{\tau^{-1}(i)}, y_i) \prod_{i=j+1}^k W_1(x_{\pi^{-1}(i)}, y_i) \\
& + (x_1 - x_{\sigma^{-1}(k)}) \sum_{\sigma} C_{\sigma}^{(k)}(x_1, \dots, y_k) W_1(x_{\sigma^{-1}(k)}, y_k) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& - \sum_{j \neq k} \sum_{\pi} \sum_{\tau} \frac{C_{\tau}^{(j)}(x_1, \dots, y_j) C_{\pi}^{(k-j)}(x_{j+1}, \dots, y_j)}{y_k - y_j} W_1(x_{\pi^{-1}(k)}, y_j) \\
& \prod_{i=1}^j W_1(x_{\tau^{-1}(i)}, y_i) \prod_{i=j+1}^{k-1} W_1(x_{\pi^{-1}(i)}, y_i) \\
& + \sum_{\sigma} \sum_{j \neq k} \frac{(x_{\sigma^{-1}(k)} - X(y_j))}{y_k - y_j} W_1(x_{\sigma^{-1}(k)}, y_j) \\
& \times \left( \operatorname{Res}_{y_k \rightarrow y_j} C_{\sigma}^{(k)} \right) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i)
\end{aligned}
\tag{4-81}$$

Using eq. (3-2) in the second line, exactly cancels the first line, and thus we get:

$$\begin{aligned}
A & = - \sum_{j \neq k} \sum_{\pi} \sum_{\tau} \frac{C_{\tau}^{(j)}(x_1, \dots, y_j) C_{\pi}^{(k-j)}(x_{j+1}, \dots, y_j)}{y_k - y_j} W_1(x_{\pi^{-1}(k)}, y_j) \\
& \prod_{i=1}^j W_1(x_{\tau^{-1}(i)}, y_i) \prod_{i=j+1}^{k-1} W_1(x_{\pi^{-1}(i)}, y_i)
\end{aligned}$$

$$\begin{aligned}
& + \sum_{\sigma} \sum_{j \neq k} \frac{(x_{\sigma^{-1}(k)} - X(y_j))}{y_k - y_j} W_1(x_{\sigma^{-1}(k)}, y_j) \\
& \quad \times \left( \operatorname{Res}_{y_k \rightarrow y_j} C_{\sigma}^{(k)} \right) \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
(4-82)
\end{aligned}$$

which is a rational fraction in  $y_k$  with poles at  $y_k = y_j$  for some  $j$ . From Lemma 4.5,  $A$  as defined in eq. (4-79) cannot have poles at  $y_k = y_j$ , thus  $A = 0$ .

Now, we have to check eq. (2-30) Using eq. (4-32), one has:

$$\begin{aligned}
& \hat{P}_k(x_1, \dots, y_k) - \hat{W}_{k-1}(x_2, \dots, x_k, y_1) \\
= & \operatorname{Pol}_{x_1} V_1'(x_1) \hat{U}_k(x_1, y_1, \dots, x_k, y_k) \\
= & \operatorname{Pol}_{x_1} Y(x_1) \sum_{\sigma \in \Sigma_k} \sum_{j \neq k} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)} \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& \quad \frac{U_1(x_{\sigma^{-1}(k)}, y_k) - U_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j} \\
= & \operatorname{Pol}_{x_1} Y(x_1) \sum_{\sigma \in \Sigma_k, \sigma(1)=k} \sum_{j \neq k} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)} \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& \quad \frac{U_1(x_1, y_k) - U_1(x_1, y_j)}{y_k - y_j} \\
& + \operatorname{Pol}_{x_1} Y(x_1) \sum_{\sigma \in \Sigma_k, \sigma(1) \neq k} \sum_{j \neq k} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)} W_1(x_1, y_{\sigma(1)}) \prod_{i \neq k, \sigma(1)} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& \quad \frac{U_1(x_{\sigma^{-1}(k)}, y_k) - U_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j} \\
= & - \sum_{\sigma \in \Sigma_k, \sigma(1)=k} \sum_{j \neq k} \sum_{l \neq 1} \operatorname{Res}_{x_1 \rightarrow x_l} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)} \prod_{i=1}^{k-1} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& \quad \frac{E(x_1, y_k) - E(x_1, y_j) - E(x_l, y_k) + E(x_l, y_j)}{(x_1 - x_l)(y_k - y_j)} \\
& + \sum_{\sigma \in \Sigma_k, \sigma(1) \neq k} \sum_{j \neq k} \sum_{l \neq 1} \operatorname{Res}_{x_1 \rightarrow x_l} \operatorname{Res}_{y'_k \rightarrow y_j} C_{\sigma}^{(k)} \prod_{i \neq k, \sigma(1)} W_1(x_{\sigma^{-1}(i)}, y_i) \\
& \quad \frac{(y_{\sigma(1)} - Y(x_1))W_1(x_1, y_{\sigma(1)}) - (y_{\sigma(1)} - Y(x_l))W_1(x_l, y_{\sigma(1)})}{x_1 - x_l} \\
& \quad \frac{U_1(x_{\sigma^{-1}(k)}, y_k) - U_1(x_{\sigma^{-1}(k)}, y_j)}{y_k - y_j} \\
(4-83)
\end{aligned}$$

In order to satisfy eq. (2-30), we must prove that  $B = 0$ , where:

$$B := \sum_{l=2}^k \frac{\hat{W}_{l-1}(x_1, y_1, \dots, y_{l-1}) - \hat{W}_{l-1}(x_l, y_1, x_2, \dots, y_{l-1})}{x_1 - x_l}$$

$$\begin{aligned}
& \times \hat{U}_{k-l+1}(x_l, y_l, \dots, x_k, y_k) \\
& + \sum_{l=1}^{k-1} \hat{W}_l(x_1, \dots, y_l) \hat{W}_{k-l}(x_k, y_l, \dots, y_{k-1}) \\
& - \hat{P}_k(x_1, y_1, x_2, \dots, x_k, y_k) \\
& - (Y(x_1) - y_k) \hat{U}_k(x_1, \dots, y_k)
\end{aligned}
\tag{4-84}$$

One does it in a way very similar to the previous one, i.e. first prove, using eq. (3-2), that  $B$  is a rational fraction of  $x_1$ , with poles at  $x_1 = x_l$ . But  $B$  can have no pole at  $x_1 = x_l$ , so,  $B = 0$ .

□

## 5 Matricial correlation functions

So far, we have computed mixed correlation functions with only one trace, i.e. the generating function of connected discrete surfaces with one boundary. In this section, we generalize this theory to the computation of generating functions of non-connected discrete surfaces with any number of boundaries. In order to derive those correlation functions, a matricial approach of the problem, similar to the one developed in [14], is used.

**Definition 5.1** *Let  $k$  be a positive integer. Let  $\pi$  and  $\pi'$  be two permutations of  $\Sigma_k$  and decompose  $\pi'^{-1}\pi$  into the product of its irreducible cycles:*

$$\pi'^{-1}\pi = P_1 P_2 \dots P_n \tag{5-1}$$

Each cycle  $P_i$  of  $\pi'^{-1}\pi$ , of length  $p_i$ , is denoted :

$$P_m = (i_{m,1} \xrightarrow{\pi} j_{m,1} \rightsquigarrow^{\pi'^{-1}} i_{m,2} \xrightarrow{\pi} j_{m,2} \rightsquigarrow^{\pi'^{-1}} \dots \rightsquigarrow^{\pi'^{-1}} i_{m,p_m} \xrightarrow{\pi} j_{m,p_m} \rightsquigarrow^{\pi'^{-1}} i_{m,1}) \tag{5-2}$$

For any  $(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \in \mathbf{C}$ , we define :

$$\mathcal{W}_{\pi, \pi'}^k(x_1, y_1, \dots, x_k, y_k) := \left\langle \prod_{m=1}^n \left( \delta_{p_m, 1} + \frac{1}{N} \text{Tr} \prod_{j=1}^{p_m} \frac{1}{(M_1 - x_{i_{m,j}})(M_2 - y_{j_{m,j}})} \right) \right\rangle \tag{5-3}$$

which is a  $k! \times k!$  matrix.

Let us now generalize the notion of planarity of a permutation.

**Definition 5.2** *Let  $k$  be a positive integer. Let  $\pi$  and  $\pi'$  be two permutations of  $\Sigma_k$ .*

A permutation  $\sigma \in \Sigma_k$  is said to be planar wrt  $(\pi, \pi')$  if

$$n_{\text{cycles}}(\pi^{-1}\sigma) + n_{\text{cycles}}(\pi'^{-1}\sigma) = k + n_{\text{cycles}}(\pi'^{-1}\pi) \quad (5-4)$$

Let  $\Sigma_k^{(\pi, \pi')} \subset \Sigma_k$  be the set of permutations planar wrt  $(\pi, \pi')$ .

Graphically, if one draws the sets of points  $(x_{i_1,1}, y_{j_1,1}, x_{i_1,2}, y_{j_1,2}, \dots, x_{i_1,p_1}, y_{j_1,p_1})$ ,  $(x_{i_2,1}, y_{j_2,1}, x_{i_2,2}, y_{j_2,2}, \dots, x_{i_2,p_2}, y_{j_2,p_2})$ ,  $\dots$ ,  $(x_{i_p,1}, y_{j_p,1}, x_{i_p,2}, y_{j_p,2}, \dots, x_{i_n,p_n}, y_{j_n,p_n})$  on  $n$  circles and link each pair  $(x_j, y_{\sigma(j)})$  by a line, these lines do not intersect nor go from one circle to another.

**Remark 5.1** One can straightforwardly see two properties of these sets:

- This relation of planarity wrt to  $(\pi, \pi')$  is symmetric in  $\pi$  and  $\pi'$ , that is to say:

$$\Sigma_k^{(\pi, \pi')} = \Sigma_k^{(\pi', \pi)} \quad (5-5)$$

- The planarity defined in 4.1 corresponds to  $\pi = Id$  and  $\pi' = S^{-1}$ :

$$\bar{\Sigma}_k = \Sigma_k^{(Id, S^{-1})} \quad (5-6)$$

Directly from these definitions and the preceding results comes the following theorem computing any generating function of discrete surface with boundaries.

**Theorem 5.1**

$$\mathcal{W}_{\pi, \pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = \sum_{\sigma} \mathcal{C}_{\sigma, \pi, \pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)})$$

(5-7)

where  $\mathcal{C}_{\sigma}^k$  is the  $k! \times k!$ -matrix defined by:

- $\mathcal{C}_{\sigma, \pi, \pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) := 0$  if  $\sigma$  is not planar wrt  $(\pi, \pi')$ ;
- if  $\sigma$  is planar wrt  $(\pi, \pi')$  :

$$\begin{aligned} \mathcal{C}_{\sigma, \pi, \pi'}^k(x_1, y_1, \dots, x_k, y_k) &:= \prod_{m=1}^a F^{(a_m)}(x_{r_{m,1}}, y_{\sigma^{-1}(r_{m,1})}, \dots, x_{r_{m,a_m}}, y_{\pi(r_{m,1})}) \\ &\quad \times \prod_{m=1}^{\tilde{a}} F^{(\tilde{a}_m)}(x_{\tilde{r}_{m,1}}, y_{\sigma(\tilde{r}_{m,1})}, \dots, x_{\tilde{r}_{m,a_m}}, y_{\pi'(\tilde{r}_{m,1})}) \end{aligned} \quad (5-8)$$

with the decompositions of  $\pi^{-1}\sigma$  and  $\pi'^{-1}\sigma$  into their products of cycles:

$$\pi^{-1}\sigma = \pi_1\pi_2 \dots \pi_a \quad , \quad \pi'^{-1}\sigma = \tilde{\pi}_1\tilde{\pi}_2 \dots \tilde{\pi}_{\tilde{a}} \quad (5-9)$$

such that:

$$\pi_m = (r_{m,1}, r_{m,2}, \dots, r_{m,a_m}) \quad , \quad \tilde{\pi}_m = (\tilde{r}_{m,1}, \tilde{r}_{m,2}, \dots, \tilde{r}_{m,\tilde{a}_m}) \quad (5-10)$$

## 5.1 Properties of the $C_\sigma^k$ 's.

**Lemma 5.1** *The matrices  $C_\sigma^k$  are symmetric.*

**proof:**

It comes directly from the definition.  $\square$

**Lemma 5.2**

$$\sum_{\sigma} C_{\sigma}^k = Id \quad (5-11)$$

**proof:**

One has:

$$\mathcal{W}_{\pi, \pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) = \sum_{\sigma} C_{\sigma, \pi, \pi'}^k \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \quad (5-12)$$

Let us shift all the  $x$ 's by a translation  $a$  and send  $a \rightarrow \infty$ , i.e. replace all the  $x_i$ 's by  $x_i + a$ . In the LHS, only the  $\delta$ -terms of def.5.1 survive in the limit  $a \rightarrow \infty$ , and thus the LHS tends towards the identity matrix. In the RHS, notice that  $W_1(x_i + a, y_{\sigma(i)}) \rightarrow 1$ . And  $C_{\sigma}^k$ , which depends only on the differences between  $x_i$ 's, is independent of  $a$ .  $\square$

**Definition 5.3**

$$\mathcal{M}^k(\vec{x}, \vec{y}, \xi, \eta)_{\pi, \pi'} := \prod_i (\delta_{\pi(i), \pi'(i)} + \frac{1}{(\xi - x_i)(\eta - y_{\pi(i)})}) \quad (5-13)$$

Let  $\mathcal{A}^{(k)}(x_1, y_1, \dots, x_k, y_k)$  be the  $k! \times k!$  defined by:

$$\begin{cases} \mathcal{A}_{\pi, \pi}^{(k)}(x_1, y_1, \dots, x_k, y_k) := \sum_i x_i y_{\pi(i)} \\ \mathcal{A}_{\pi, \pi'}^{(k)}(x_1, y_1, \dots, x_k, y_k) := 1 \text{ if } \pi \pi'^{-1} = \text{transposition} \\ \mathcal{A}_{\pi, \pi'}^{(k)}(x_1, y_1, \dots, x_k, y_k) := 0 \text{ otherwise} \end{cases} \quad (5-14)$$

**Theorem 5.2**

$$\boxed{\forall \sigma, \xi, \eta, \quad [\mathcal{M}^k(\vec{x}, \vec{y}, \xi, \eta), C_{\sigma}^k(\vec{x}, \vec{y})] = 0} \quad (5-15)$$

and

$$\boxed{\forall \xi, \eta, \quad [\mathcal{M}^k(\vec{x}, \vec{y}, \xi, \eta), \mathcal{W}^k(\vec{x}, \vec{y})] = 0} \quad (5-16)$$

**Corollary 5.1**

$$\forall \sigma, \quad [\mathcal{A}^{(k)}(\vec{x}, \vec{y}), C_{\sigma}^k(\vec{x}, \vec{y})] = 0 \quad (5-17)$$



Theorem 5.2 is a very powerful equation. One can check on small values of  $k$ , that theorem 5.2, as well as

$$\mathcal{C}_{\sigma,\pi,\pi}^k(\vec{x}, \vec{y}) = \delta_{\sigma,\pi} \quad (5-18)$$

is sufficient to determine the matrix  $\mathcal{C}_{\sigma}^k(\vec{x}, \vec{y})$ , and thus  $\mathcal{W}^k(\vec{x}, \vec{y})$ .

**proof:**

Let us define:

$$\tilde{\mathcal{M}}(\vec{x}, \vec{y}, \xi, \eta) := \mathcal{M}(N\vec{x}, \vec{y}, N\xi, \eta) \quad (5-19)$$

and

$$\tilde{\mathcal{W}}_{\pi,\pi'}^k(x_1, y_1, \dots, x_k, y_k) := \left\langle \prod_{m=1}^n \left( \delta_{p_m,1} + \text{Tr} \prod_{j=1}^{p_m} \frac{1}{N} \frac{1}{(M_1 - x_{i_m,j})(M_2 - y_{j_m,j})} \right) \right\rangle \quad (5-20)$$

It was proven in [14] that:

$$[\tilde{\mathcal{M}}^k(\vec{x}, \vec{y}, \xi, \eta), \tilde{\mathcal{W}}^k(\vec{x}, \vec{y})] = 0 \quad (5-21)$$

Now, in the large  $N$  limit, the factorization property [5]  $\langle \text{Tr Tr} \rangle \sim \langle \text{Tr} \rangle \langle \text{Tr} \rangle$ , implies:

$$\begin{aligned} & \tilde{\mathcal{W}}_{\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \\ & \sim N^{n_{\text{cycles}}(\pi'^{-1}\pi) - k} \prod_{m=1}^n W_{p_m}(x_{i_m,1}, y_{\pi(i_m,1)}, \dots, x_{i_m,p_m}, y_{\pi(i_m,p_m)}) \\ & \sim N^{n_{\text{cycles}}(\pi'^{-1}\pi) - k} \mathcal{W}_{\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \end{aligned} \quad (5-22)$$

and using theorem 5.1, we have:

$$\begin{aligned} & \tilde{\mathcal{W}}_{\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \\ & \sim N^{n_{\text{cycles}}(\pi'^{-1}\pi) - k} \sum_{\sigma} \mathcal{C}_{\sigma,\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \end{aligned} \quad (5-23)$$

Notice that

$$\begin{aligned} & \mathcal{C}_{\sigma,\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \\ & = N^{k - n_{\text{cycles}}(\pi^{-1}\sigma) + k - n_{\text{cycles}}(\pi'^{-1}\sigma)} \mathcal{C}_{\sigma,\pi,\pi'}^k(Nx_1, y_1, Nx_2, y_2, \dots, Nx_k, y_k) \\ & = N^{k - n_{\text{cycles}}(\pi'^{-1}\pi)} \mathcal{C}_{\sigma,\pi,\pi'}^k(Nx_1, y_1, Nx_2, y_2, \dots, Nx_k, y_k) \end{aligned} \quad (5-24)$$

Thus:

$$\tilde{\mathcal{W}}_{\pi,\pi'}^k(x_1, y_1, x_2, y_2, \dots, x_k, y_k)$$

$$\sim \sum_{\sigma} \mathcal{C}_{\sigma, \pi, \pi'}^k(Nx_1, y_1, Nx_2, y_2, \dots, Nx_k, y_k) \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \quad (5-25)$$

Then, from [14], we have:

$$0 = \sum_{\sigma} [\mathcal{M}^k(N\vec{x}, \vec{y}, N\xi, \eta), \mathcal{C}_{\sigma}^k(Nx_1, y_1, \dots, Nx_k, y_k)] \prod_{i=1}^k W_1(x_i, y_{\sigma(i)}) \quad (5-26)$$

In particular, choose a permutation  $\sigma$ , and take the limit where  $y_i \rightarrow Y(x_{\sigma^{-1}(i)})$ , you get in that limit:

$$0 = \left[ \mathcal{M}^k(N\vec{x}, Y(x_{\sigma^{-1}}), N\xi, \eta), \mathcal{C}_{\sigma}^k(Nx_1, Y(x_{\sigma^{-1}(1)}), \dots, Nx_k, Y(x_{\sigma^{-1}(k)})) \right] \quad (5-27)$$

Since this equation holds for any potentials  $V_1$  and  $V_2$ , it holds for any function  $Y(x)$ , and thus the  $Y(x_i)$ 's can be chosen independently of the  $x_i$ 's, and thus, for any  $y_1, \dots, y_k$ , we have:

$$0 = [\mathcal{M}^k(N\vec{x}, \vec{y}, N\xi, \eta), \mathcal{C}_{\sigma}^k(Nx_1, y_1, \dots, Nx_k, y_k)] \quad (5-28)$$

Since it holds for any  $x_i$ 's and  $\xi$ , it also holds for  $x_i/N$  and  $\xi/N$ .  $\square$

The corollary is obtained by taking the large  $\xi$  and  $\eta$  limit.  $\square$

## 5.2 Examples: $k = 2$ .

$$\mathcal{W}^{(2)} = \begin{pmatrix} W_{11}W_{22} & \frac{W_{11}W_{22}-W_{12}W_{21}}{(x_1-x_2)(y_1-y_2)} \\ \frac{W_{11}W_{22}-W_{12}W_{21}}{(x_1-x_2)(y_1-y_2)} & W_{12}W_{21} \end{pmatrix} \quad (5-29)$$

where  $W_{ij} = W_1(x_i, y_j)$ .

$$\mathcal{C}_{Id}^2 = \begin{pmatrix} 1 & \frac{1}{(x_1-x_2)(y_1-y_2)} \\ \frac{1}{(x_1-x_2)(y_1-y_2)} & 0 \end{pmatrix} \quad (5-30)$$

$$\mathcal{C}_{(12)}^2 = \begin{pmatrix} 0 & \frac{1}{(x_1-x_2)(y_2-y_1)} \\ \frac{1}{(x_1-x_2)(y_2-y_1)} & 1 \end{pmatrix} = 1 - \mathcal{C}_{Id}^2 \quad (5-31)$$

## 6 Gaussian case

There is an example of special interest, in particular for its applications to string theory [16], it is the gaussian-complex matrix model case,  $V_1 = V_2 = 0$ . In that case one has  $E(x, y) = xy - 1$ , and thus:

$$W_1(x, y) = \frac{xy}{xy - 1} \quad (6-1)$$

The loop equation defining recursively the  $W_k$ 's can be written:

$$(x_1 y_k - 1)W_k(x_1, y_1, \dots, x_k, y_k) = x_1 \sum_j \frac{W_{j-1}(x_j, y_1, \dots, y_{j-1}) - W_{j-1}(x_1, y_1, \dots, y_{j-1})}{x_1 - x_j} \times W_{k-j+1}(x_j, y_j, \dots, x_k, y_k) \quad (6-2)$$

Its solution is then:

$$W_k(x_1, y_1, \dots, x_k, y_k) = \sum_{\sigma \in \Sigma_k} C_\sigma^{(k)}(x_1, y_1, \dots, x_k, y_k) \prod_{i=1}^k \frac{x_i y_{\sigma(i)}}{x_i y_{\sigma(i)} - 1} \quad (6-3)$$

From the loop equation, one can see that  $W_k(x_1, y_1, \dots, x_k, y_k)$  may have poles only when  $x_i \rightarrow y_j$  for any  $i$  and  $j$ . Because the  $C_\sigma$ 's are rational functions of all these variables, one can write:

$$W_k(x_1, y_1, \dots, x_k, y_k) = \frac{N_k(x_1, y_1, x_2, y_2, \dots, x_k, y_k)}{\prod_{i,j} (x_i y_j - 1)} \quad (6-4)$$

where  $N_k(x_1, y_1, x_2, y_2, \dots, x_k, y_k)$  is a polynomial in all its variables.

Moreover, the loop equation taken for the values  $x_k = 0$  or  $y_k = 0$  shows that  $W_k(x_1, y_1, \dots, 0, y_k) = W_k(x_1, y_1, \dots, x_k, 0) = 0$ . Using the cyclicity property of  $W_k(x_1, y_1, \dots, x_k, y_k)$ , one can claim that it vanishes whenever one of its arguments is equal to 0. One can thus factorize the polynomial  $N(x_1, y_1, x_2, y_2, \dots, x_k, y_k)$  as follows:

$$W_k(x_1, y_1, \dots, x_k, y_k) = \frac{Q_k(x_1, y_1, x_2, y_2, \dots, x_k, y_k) \prod_i x_i y_i}{\prod_{i,j} (x_i y_j - 1)} \quad (6-5)$$

where  $Q(x_1, y_1, \dots, x_k, y_k)$  is a polynomial itself.

As an example, we have:

- for  $k = 2$ :

$$W_2(x_1, y_1, x_2, y_2) = \frac{x_1 x_2 y_1 y_2}{\prod_{i,j} (x_i y_j - 1)} \quad \text{and} \quad Q_2(x_1, y_1, x_2, y_2) = 1 \quad (6-6)$$

- for  $k = 3$ :

$$W_3(x_1, y_1, x_2, y_2, x_3, y_3) = (2 - \sum_i x_i y_{i+1} + x_1 x_2 x_3 y_1 y_2 y_3) \frac{x_1 x_2 x_3 y_1 y_2 y_3}{\prod_{i,j} (x_i y_j - 1)} \quad (6-7)$$

and

$$Q_3(x_1, y_1, x_2, y_2, x_3, y_3) = (2 - x_1 y_2 - x_2 y_3 - x_3 y_1 + x_1 x_2 x_3 y_1 y_2 y_3) \quad (6-8)$$

## 7 Conclusion

In this article, we have computed the generating functions of discs with all possible boundary conditions, i.e. the large  $N$  limit of all correlation functions of the formal 2-matrix model. We have found that the  $2k$  point correlation function, can be written like the Bethe ansatz for the  $\delta$ -interacting bosons, i.e. a sum over permutations of product of 2-point functions. That formula is universal, it is independent of the potentials.

An even more powerful approach consists in gathering all possible  $2k$  point correlation functions in a  $k! \times k!$  matrix  $\mathcal{W}^k$ . We have found that this matrix  $\mathcal{W}^k$  satisfies commutation relations with a family of matrices  $\mathcal{M}^k$  which depend on two spectral parameters, and are related to the representations of  $U(n)$  [14]. We claim that the theorem 5.2 is almost equivalent to the loop equations, and allows to determine  $\mathcal{W}^k$ .

It remains to understand how all these matrices and coefficients  $C_\sigma$  are related to usual formulations of integrability, i.e. how to write these in terms of Yang Baxter equations.

One could also hope to find a direct proof of theorem 3.2, without having to solve the loop equations. In other words, we have found that the  $2k$ -point function can be written only in terms of  $W_1$ , while, in the derivation, we use the one point functions  $Y(x)$  and  $X(y)$  although they don't appear in the final result.

The next step, is to be able to compute the  $1/N^2$  expansion of those correlation functions, as well as the large  $N$  limit of connected correlation functions. We are already working that out, by mixing the approach presented in the present article, and the Feynman graph approach of [8], generalized to the 2-matrix model in [9].

Another prospect, is to go to the critical limit, i.e. where we describe generating function for continuous surfaces with conformal invariance, and interpret this as boundary conformal field theory [].

## Acknowledgements

The authors want to thank M. Bauer, M. Bergère, F. David, P. Di Francesco, J.B. Zuber for stimulating discussions. This work was partly supported by the european network Enigma (MRTN-CT-2004-5652).

## References

- [1] M. Bergère, "Biorthogonal Polynomials for Potentials of two Variables and External Sources at the Denominator", hep-th/0404126, SPhT-T04/042.

- [2] M. Bertola, B. Eynard, “Mixed Correlation Functions of the Two-Matrix Model”, *J. Phys. A* **36** (2003) 7733-7750, xxx, hep-th/0303161.
- [3] G. Bonnet, F. David, B. Eynard, “Breakdown of universality in multi-cut matrix models”, *J. Phys. A* **33** 6739-6768 (2000).
- [4] E. Brezin, C. Itzykson, G. Parisi, and J. Zuber, *Comm. Math. Phys.* **59**, 35 (1978).
- [5] P. Di Francesco, P. Ginsparg, J. Zinn-Justin, “2D Gravity and Random Matrices”, *Phys. Rep.* **254**, 1 (1995).
- [6] B. Eynard, “Master loop equations, free energy and correlations for the chain of matrices”, *JHEP*11(2003)018, xxx, hep-th/0309036, ccscd-00000572.
- [7] B. Eynard, “Large N expansion of the 2-matrix model”, *JHEP* **01** (2003) 051, hep-th/0210047.
- [8] B. Eynard, “Topological expansion for the 1-hermitian matrix model correlation functions”, *JHEP*JHEP/024A/0904, xxx, hep-th/0407261.
- [9] B. Eynard, N. Orantin, “A residue formula for correlation functions for the hermitian 2-matrix model”, in preparation.
- [10] B. Eynard, “Eigenvalue distribution of large random matrices, from one matrix to several coupled matrices” *Nucl. Phys. B***506**, 633 (1997), cond-mat/9707005.
- [11] B. Eynard, “Correlation functions of eigenvalues of multi-matrix models, and the limit of a time dependent matrix”, *J. Phys. A: Math. Gen.* **31**, 8081 (1998), cond-mat/9801075.
- [12] B. Eynard “An introduction to random matrices”, lectures given at Saclay, October 2000, notes available at <http://www-spht.cea.fr/articles/t01/014/>.
- [13] B. Eynard “Polynômes biorthogonaux, problème de Riemann–Hilbert et géométrie algébrique”, Habilitation à diriger les recherches, université Paris VII, (2005).
- [14] B. Eynard, A. Prats-Ferrer, “2-matrix versus complex matrix model, integrals over the unitary group as triangular integrals”, preprint SPHT T05/011, preprint UB-ECM-PF 05/03, xxx, hep-th/0502041.
- [15] B. Eynard, M.L. Mehta, “Matrices coupled in a chain: eigenvalue correlations”, *J. Phys. A: Math. Gen.* **31**, 4449 (1998), cond-mat/9710230.

- [16] C. Kristjansen, J. Plefka, G. W. Semenoff, M. Staudacher, “A New Double-Scaling Limit of N=4 Super Yang-Mills Theory and PP-Wave Strings”, *Nucl.Phys.* **B643** (2002) 3-30, hep-th/0205033.
- [17] M. Gaudin M “La Fonction d’Onde de Bethe”, Collection du Commissariat a l’Energie Atomique: Serie, Editor: Masson, (1983).
- [18] V.A. Kazakov, “Ising model on a dynamical planar random lattice: exact solution”, *Phys Lett.* **A119**, 140-144 (1986).
- [19] V. Kazakov, I. Kostov, ”Loop gas model for open strings”, *Nuc. Phys.* **B386** (1992) 520.
- [20] V.A. Kazakov, A. Marshakov, ”Complex Curve of the Two Matrix Model and its Tau-function”, *J.Phys.* **A36** (2003) 3107-3136, hep-th/0211236.
- [21] I. K. Kostov, “Boundary Correlators in 2D Quantum Gravity: Liouville versus Discrete Approach”, *Nucl.Phys.* **B658** (2003) 397-416, hep-th/0212194.
- [22] A. Matytsin, “on the large  $N$  limit of the Itzykson Zuber Integral”, *Nuc. Phys.* **B411**, 805 (1994), hep-th/9306077.
- [23] M.L. Mehta, *Random Matrices*, 2nd edition, (Academic Press, New York, 1991).
- [24] M.L. Mehta, ”A method of integration over matrix variables”, *Comm. Math. Phys.* **bf 79** (1981) 327-340.
- [25] M. Staudacher, “Combinatorial solution of the 2-matrix model”, *Phys. Lett.* **B305** (1993) 332-338.