

SURPRISING SIMPLICITY OF $\mathcal{N} = 8$ SUPERGRAVITY*

N. E. J. BJERRUM-BOHR

*The Niels Bohr International Academy, The Niels Bohr Institute,
Blegdamsvej 17, DK-2100, Copenhagen Ø, Denmark
bjbohr@nbi.dk*

PIERRE VANHOVE

*Institut des Hautes Etudes Scientifiques, Le Bois-Marie,
F-91440 Bures-sur-Yvette, France
CEA, DSM, Institut de Physique Théorique, IPhT, CNRS, MPPU,
URA2306, Saclay, F-91191 Gif-sur-Yvette, France
pierre.vanhove@cea.fr*

Communicated by D. V. Ahluwalia

Gravity amplitudes are, via the Kawai–Lewellen–Tye relations intimately linked to products of Yang–Mills amplitudes. Explicitly, this shows up in computations of $\mathcal{N} = 8$ supergravity where the perturbative expansion and ultraviolet behavior of this theory is akin to $\mathcal{N} = 4$ super-Yang–Mills at least through three loops. Full persistence to all loop orders would be truly remarkable and imply finiteness of $\mathcal{N} = 8$ supergravity in four dimensions.

1. Introduction

Since the discovery of quantum mechanics in the previous century, physicists have been pursuing the construction of a fundamental theory for quantum gravity. Quantum-gravitational effects appear to be essential in understanding the physics of very dense matter objects such as the early universe and black holes. However, although the searches for a theory of quantum gravity have been diverse, extensive and many, the fundamental concepts of such a theory are still elusive. General relativity provides us with a very successful theory for gravity which captures the apparent necessary knowledge for a complete treatment of the gravitational attraction and its intimate connection with matter, space and time. However, it is incompatible with basic quantum-mechanical ideas such as operator space and expectation values. A traditional approach to perturbative gravity through a Lagrangian description is possible although complicated by a divergent ultraviolet behavior. Progress

*This essay received an honorable mention in the 2009 Essay Competition of the Gravity Research Foundation. It was refereed, not as a regular IJMPD research paper, but as an essay.

has, however, been achieved this way through treating gravity as an effective field theory.^{1–5}

For many years the combination of supersymmetry with a Lagrangian description of quantum gravity was considered to be a way out of the troublesome ultraviolet-divergent behavior of such a theory due to the introduction of extra fundamental symmetries. Such theories were termed *supergravity models*. The possibly most famous one is the model of maximal $\mathcal{N} = 8$ supergravity.^{6–8} However, with the advent of superstring theory in the mid-1980s such models were abandoned due a common belief of unavoidable divergences in their perturbative expansion via power-counting arguments,^{9–12} relegating these theories to low energy effective descriptions of string theory.

In recent years, due to remarkable progress in computational techniques by combining various inputs from string theory, extended supersymmetry and unitarity, there has been a renewed interest in supergravity models for quantum gravity and it has become clear that $\mathcal{N} = 8$ supergravity has a much better perturbative expansion than power-counting naïvely predicts. Surprisingly, the ultraviolet behavior of $\mathcal{N} = 8$ supergravity occurs explicitly to be identical to that of $\mathcal{N} = 4$ super-Yang–Mills to at least three loops^{13–20} — very likely six loops.^{21,22} If this identical UV behavior persists to all orders in perturbation theory, then $\mathcal{N} = 8$ supergravity will be ultraviolet-finite in four dimensions.²¹

The massless spectrum of $\mathcal{N} = 8$ supergravity can be seen as the tensorial product of two copies of $\mathcal{N} = 4$ super-Yang–Mills theories, through the Kawai–Lewellen–Tye relations,²³ which are motivated by string theory. In these relations the massless supergravity (closed string) vertex operators are written as the left/right product of Yang–Mills open string vertex operators. One can hence organize $\mathcal{N} = 8$ supergravity tree-level amplitudes according to a relation^{13,24–31} which we will write schematically, in the following way:

$$\text{Gravity} \sim (\text{Yang–Mills}) \times (\text{Yang–Mills}'). \quad (1.1)$$

This simple relation between a theory of gravity and two gauge theories is observed directly in on-shell S matrix elements but appears to be rather odd at the level of the Lagrangian and its interactions. (This is true even if part of the Lagrangian is rearranged as a product of Yang–Mills types of interactions at the two-derivative level^{32,33} or for higher derivative corrections.³⁴)

2. Supergravity Amplitudes

A superficial power-counting argument indicates that an L -loop n -graviton amplitude in D dimensions behaves as

$$[\mathfrak{M}_L^{(D)}] = \text{mass}^{(D-2)L+2}. \quad (2.1)$$

This count can be compared to the superficial power-counting of the four-gluon amplitude in $\mathcal{N} = 4$ super-Yang–Mills, which is given by

$$[\mathfrak{A}_{4;L}^{(D)}] = \text{mass}^{(D-4)L}. \quad (2.2)$$

For $\mathcal{N} = 4$ super-Yang–Mills in four dimensions, we see that the theory is at most logarithmically diverging (since the coupling constant is dimensionless). The extended $\mathcal{N} = 4$ supersymmetry guarantees perturbative finiteness.^{35,36} Color-ordered amplitudes factorize the dimension 4 operator F^4 at one-loop order and the dimension 6 operator $\partial^2 F^4$ at higher-loop order and hence satisfy the dimensional analysis

$$[\mathfrak{A}_{4;L}^{(D)}] = \text{mass}^{(D-4)L-6}[\partial^2 F^4]. \tag{2.3}$$

This implies that L -loop four-point amplitudes in $\mathcal{N} = 4$ super-Yang–Mills are ultraviolet-divergent in dimensions

$$D \geq 4 + \frac{6}{L}, \tag{2.4}$$

thus implying perturbative ultraviolet finiteness in $D = 4$ dimensions (the negative mass dimension reflects the infrared behavior of the amplitude).

The difference between the formulae (2.1) and (2.3) reflects the difference in dimensions of the coupling constant of the two theories. However, this superficial power-counting misses dramatic simplifications taking place in on-shell amplitudes due to the extended $\mathcal{N} = 8$ supersymmetry²² and the role of (diffeomorphism) gauge invariance.^{17,37}

String-based methods for constructing higher-loop amplitudes indicate^{21,22} that the perturbative behavior of $\mathcal{N} = 8$ supergravity amplitudes is improved by the factorization of the dimension 8 R^4 operator together with extra powers of derivatives

$$[\mathfrak{M}_L^{(D)}] = \text{mass}^{(D-2)L-6-2\beta_L}[\partial^{2\beta_L} R^4], \tag{2.5}$$

with the $\beta_L = L$ rule:

$$\beta_1 = 0; \quad \beta_L = L \quad \text{for } 2 \leq L. \tag{2.6}$$

This leads to a superficial ultraviolet behavior for $\mathcal{N} = 8$ supergravity amplitudes of

$$[\mathfrak{M}_L^{(D)}] = \text{mass}^{(D-4)L-6}[D^{2L} R^4], \tag{2.7}$$

which is *similar* to the ultraviolet behavior in (2.3) for $\mathcal{N} = 4$ super-Yang–Mills. When the $\beta_L = L$ rule (2.6) is satisfied, the $\mathcal{N} = 8$ four-graviton supergravity amplitude has the same critical dimension (2.4) for ultraviolet divergences as $\mathcal{N} = 4$ super-Yang–Mills.

The validity of the $\beta_L = L$ rule to all orders in perturbation theory implies perturbative finiteness of the four-graviton $\mathcal{N} = 8$ supergravity amplitude in four dimensions.

3. $\mathcal{N} = 8$ Supergravity as a Product of $\mathcal{N} = 4$ Yang–Mills

On-shell recursion relations provide very simple means of constructing $\mathcal{N} = 8$ supergravity tree-level amplitudes from three-point vertices.^{38,39} Gravity three-point vertices are given directly as squares of $\mathcal{N} = 4$ super-Yang–Mills vertices. Thus, the

resulting massless n -point tree-level amplitudes can be presented in a form involving terms with sums of squares of three-point $\mathcal{N} = 4$ super-Yang–Mills vertices.³¹

At the field theory level this amounts to replacing the gauge degree of freedom of the Yang–Mills fields by Lorentz degrees of freedom as follows:

$$A_\mu^a \rightarrow \zeta_\mu^a. \quad (3.1)$$

Such a correspondence is compatible with extended supersymmetry and can be used naïvely to promote $\mathcal{N} = 4$ super-Yang–Mills invariants to higher-derivative $\mathcal{N} = 8$ superinvariants. However, it is not possible to capture all of these by such a map,³⁴ since the full diffeomorphism invariance carries even more symmetry (such as Ricci cycling identities $R_{\mu[\nu\rho\sigma]} = 0$). The naïve application of the above substitution rule would, for instance, lead to an $\mathcal{N} = 8$ supergravity amplitude with an apparent factorization of the operator $D^2 R^4$. This would consequently make the $L = 2$ two-loop four-graviton amplitude diverge to $D = 6$. However, this is contrary to explicit knowledge, since this amplitude has been shown to be finite up to $D \leq 6$.¹³

The main reason for this glitch is that diffeomorphism invariance for $\mathcal{N} = 8$ supergravity implies that one must sum over all permutations of external legs. Doing this makes the invariant $D^2 R^4 \sim (s + t + u)R^4 = 0$ vanish by on-shell momentum conservation. Consequently the first nonvanishing contribution is of order $D^4 R^4$. At two-loop order the operator $D^4 R^4$ satisfies the rule $\beta_2 = 2$ of Eq. (2.6), providing the suitable structure for the two-loop amplitude kinematic factor. However, this does not give the correct contribution for the higher-loop amplitudes which have $\beta_L \geq 3$ for $L \geq 3$.^{18,21}

Thus, we see that one has to be careful with such arguments since gravity theories have symmetries which are beyond what is provided via two copies of the gauge transformations of Yang–Mills theories.

An important consequence of the full crossing symmetry provided via the absence of the concept of color in gravity theories is that infrared divergences in quantum gravity can be treated as in QED and are much milder than in color-ordered theories like QCD. One benefit of the structure of gravitational interactions is that there are no divergences for the emission of a soft graviton from a hard line, contrary to massless QED.⁴⁰ This indicates that although gravitational interactions looks much more complicated than gauge theory ones, important simplification occurs in on-shell amplitudes at tree level. The fact that gravity amplitudes are unordered implies the no-triangle property of $\mathcal{N} = 8$ supergravity loop amplitudes^{16,17} and puts nontrivial constraints on the structure of higher-loop amplitudes.⁴¹

4. Vacuum Structure and E_7 Invariance

The $\beta_L = L$ rule can be derived up to six loops from the zero mode sector of the pure spinor formalism⁴² for four-graviton amplitudes. This shows that the role

of extended supersymmetry in perturbative $\mathcal{N} = 8$ supergravity is beyond the superspace transformation properties of the product of two $\mathcal{N} = 4$ super-Yang–Mills theories. Further analysis shows that the vacuum structure of $\mathcal{N} = 4$ super-Yang–Mills and of $\mathcal{N} = 8$ supergravity theory is very different. Mathematically, in four dimensions the vacuum of $\mathcal{N} = 8$ supergravity can be described by the homogeneous space of $\mathcal{M} = E_{7(7)}/[\text{SU}(8)_R/\mathbb{Z}_2]$. While the local symmetry group $\text{SU}(8)_R$ transforms as a “square” of the group $\text{SU}(4)_R$ (corresponding to each $\mathcal{N} = 4$ super-Yang–Mills theory), there is no concept of the symmetry of the global group $E_{7(7)}$ in $\mathcal{N} = 4$ super-Yang–Mills. The global $E_{7(7)}$ symmetry does put severe enough constraints on counterterms $\mathcal{N} = 8$ in supergravity to possibly protect the theory from diverging before nine loops, and in conjunction with the full crossing symmetry this could be enough to imply finiteness of the theory in four dimensions. As a global symmetry rotating the different vacua of $\mathcal{N} = 8$ supergravity, the $E_{7(7)}$ symmetry relates the perturbative contributions to the nonperturbative black hole production at high energy, which are required for a consistent definition of the theory.⁴³

5. Discussion

Although we are still searching for a fundamental theory of quantum gravity, we are in these years gaining a much better understanding of the necessary concepts for the formulation of such a theory. The role of dualities in supergravity theories is important for their quantization, and such investigations provide a framework for gathering further knowledge about quantum gravity, its fundamental degrees of freedom and its relation to gauge theory. A clear understanding of the question of ultraviolet finiteness and the validity of the $\beta_L = L$ rule²¹ of $\mathcal{N} = 8$ supergravity would indeed be remarkable and have huge implications for non-supersymmetric low energy descriptions of quantum gravity theories.

Acknowledgments

N. E. J. Bjerrum-Bohr is Knud Højgaard Assistant Professor at the Niels Bohr International Academy. This research was supported in part (P. V.) by the ANR grant BLAN06-3-137168.

References

1. J. F. Donoghue, *Phys. Rev. D* **50** (1994) 3874 [arXiv:gr-qc/9405057].
2. N. E. J. Bjerrum-Bohr, *Phys. Rev. D* **66** (2002) 084023 [arXiv:hep-th/0206236].
3. N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, *Phys. Rev. D* **68** (2003) 084005 [Erratum *Phys. Rev. D* **71** (2005) 069904] [arXiv:hep-th/0211071].
4. N. E. J. Bjerrum-Bohr, J. F. Donoghue and B. R. Holstein, *Phys. Rev. D* **67** (2003) 084033 [Erratum *Phys. Rev. D* **71** (2005) 069903] [arXiv:hep-th/0211072].
5. B. R. Holstein and A. Ross, arXiv:0802.0716 (hep-ph).
6. E. Cremmer, B. Julia and J. Scherk, *Phys. Lett. B* **76** (1978) 409.

7. E. Cremmer and B. Julia, *Phys. Lett. B* **80** (1978) 48.
8. E. Cremmer and B. Julia, *Nucl. Phys. B* **159** (1979) 141.
9. P. S. Howe and U. Lindstrom, *Nucl. Phys. B* **181** (1981) 487.
10. R. E. Kallosh, *Phys. Lett. B* **99** (1981) 122.
11. P. S. Howe, K. S. Stelle and P. K. Townsend, *Nucl. Phys. B* **236** (1984) 125.
12. P. S. Howe and K. S. Stelle, *Phys. Lett. B* **554** (2003) 190 [arXiv:hep-th/0211279].
13. Z. Bern, L. J. Dixon, D. C. Dunbar, M. Perelstein and J. S. Rozowsky, *Nucl. Phys. B* **530** (1998) 401 [arXiv:hep-th/9802162].
14. Z. Bern, N. E. J. Bjerrum-Bohr and D. C. Dunbar, *J. High Energy Phys.* **0505** (2005) 056 [arXiv:hep-th/0501137].
15. N. E. J. Bjerrum-Bohr, D. C. Dunbar and H. Ita, *Phys. Lett. B* **621** (2005) 183 [arXiv:hep-th/0503102].
16. N. E. J. Bjerrum-Bohr, D. C. Dunbar, H. Ita, W. B. Perkins and K. Risager, *J. High Energy Phys.* **0612** (2006) 072 [arXiv:hep-th/0610043].
17. N. E. J. Bjerrum-Bohr and P. Vanhove, *J. High Energy Phys.* **0804** (2008) 065 [arXiv:0802.0868 (hep-th)].
18. Z. Bern, J. J. Carrasco, L. J. Dixon, H. Johansson, D. A. Kosower and R. Roiban, *Phys. Rev. Lett.* **98** (2007) 161303 [arXiv:hep-th/0702112].
19. Z. Bern, J. J. M. Carrasco, L. J. Dixon, H. Johansson and R. Roiban, *Phys. Rev. D* **78** (2008) 105019 [arXiv:0808.4112 (hep-th)].
20. Z. Bern, J. J. M. Carrasco and H. Johansson, arXiv:0902.3765 (hep-th).
21. M. B. Green, J. G. Russo and P. Vanhove, *J. High Energy Phys.* **0702** (2007) 099.
22. M. B. Green, J. G. Russo and P. Vanhove, *Phys. Rev. Lett.* **98** (2007) 131602 [arXiv:hep-th/0611273].
23. H. Kawai, D. C. Lewellen and S. H. H. Tye, *Nucl. Phys. B* **269** (1986) 1.
24. Z. Bern and A. K. Grant, *Phys. Lett. B* **457** (1999) 23 [arXiv:hep-th/9904026].
25. S. Ananth and S. Theisen, *Phys. Lett. B* **652** (2007) 128 [arXiv:0706.1778 (hep-th)].
26. Z. Bern, *Living Rev. Rel.* **5** (2002) 5 [arXiv:gr-qc/0206071].
27. Z. Bern, A. De Freitas and H. L. Wong, *Phys. Rev. Lett.* **84** (2000) 3531 [arXiv:hep-th/9912033].
28. N. E. J. Bjerrum-Bohr, *Phys. Lett. B* **560** (2003) 98 [arXiv:hep-th/0302131].
29. N. E. J. Bjerrum-Bohr, *Nucl. Phys. B* **673** (2003) 41 [arXiv:hep-th/0305062].
30. N. E. J. Bjerrum-Bohr and K. Risager, *Phys. Rev. D* **70** (2004) 086011 [arXiv:hep-th/0407085].
31. Z. Bern, J. J. M. Carrasco and H. Johansson, *Phys. Rev. D* **78** (2008) 085011 [arXiv:0805.3993 (hep-ph)].
32. S. Ananth and S. Theisen, *Phys. Lett. B* **652** (2007) 128 [arXiv:0706.1778 (hep-th)].
33. S. Ananth, arXiv:0902.3128 (hep-th).
34. K. Peeters, P. Vanhove and A. Westerberg, *Class. Quant. Grav.* **18** (2001) 843 [arXiv:hep-th/0010167].
35. S. Mandelstam, *Nucl. Phys. B* **213** (1983) 149.
36. G. Bossard, P. S. Howe and K. S. Stelle, arXiv:0901.4661 (hep-th).
37. S. Badger, N. E. J. Bjerrum-Bohr and P. Vanhove, *J. High Energy Phys.* **0902** (2009) 038 [arXiv:0811.3405 (hep-th)].
38. R. Britto, F. Cachazo and B. Feng, *Nucl. Phys. B* **725** (2005) 275 [arXiv:hep-th/0412103].
39. R. Britto, F. Cachazo, B. Feng and E. Witten, *Phys. Rev. Lett.* **94** (2005) 181602 [arXiv:hep-th/0501052].

40. S. Weinberg, *Phys. Rev.* **140** (1965) B516.
41. Z. Bern, L. J. Dixon and R. Roiban, *Phys. Lett. B* **644** (2007) 265 [arXiv:hep-th/0611086].
42. N. Berkovits, *Phys. Rev. Lett.* **98** (2007) 211601 [arXiv:hep-th/0609006].
43. M. B. Green, H. Ooguri and J. H. Schwarz, *Phys. Rev. Lett.* **99** (2007) 041601.