

Tracy-Widom GUE law and symplectic invariants

*G. Borot*¹, *B. Eynard*²

¹ Institut de Physique Théorique de Saclay,
F-91191 Gif-sur-Yvette Cedex, France

² CERN, Theory Division
CH-1211 Geneva 23, Switzerland

Abstract

We establish the relation between two objects: an integrable system related to Painlevé II equation, and the symplectic invariants of a certain plane curve Σ_{TW} describing the average eigenvalue density of a random hermitian matrix spectrum near a hard edge (a bound for its maximal eigenvalue). This explains directly how the Tracy-Widom law F_{GUE} , governing the distribution of the maximal eigenvalue in hermitian random matrices, can also be recovered from symplectic invariants.

¹gaetan.borot@cea.fr

²bertrand.eynard@cea.fr

Contents

1	Basics around the topological recursion	5
1.1	Definitions	5
1.2	Matrix models and topological recursion	6
1.3	2×2 differential system and topological recursion	7
1.4	Generalities on the Schrödinger equation	9
1.5	Tau function and topological recursion	10
1.5.1	Definition of the τ -function	10
1.5.2	Large N limit	11
1.5.3	$1/N$ expansion of the τ -function	12
2	Refinement on the assumptions	13
2.1	A remark on hypothesis (iii)	13
2.2	Loop insertion operator and properties	14
2.3	Analyticity properties of the $1/N$ expansion	17
2.3.1	Summary of the problem	17
2.3.2	Analytical properties	18
2.4	Conclusion	19
3	Application to the Tracy-Widom GUE law	20
3.1	Tracy-Widom GUE and integrability	20
3.2	$s \rightarrow -\infty$ asymptotics and spectral curve	21
3.3	Existence of the $1/N$ expansion	23
3.4	Tau-function and symplectic invariants	25
3.5	Relation with the 1-matrix model with hard edge	26
3.6	Remark on correlation functions	28
4	Conclusion	29
A	A counterexample to the $1/N^2$ expansion	31
B	Topological recursion and symplectic invariants	32

Motivation

The famous Tracy-Widom law [32], describes the statistics of the largest eigenvalue of a random hermitian matrix of the GUE ensemble [28].

Consider the large deviation function

$$\mathcal{F}_N(a) \equiv \ln \text{Prob}[\lambda_{\max} \leq a] \quad \text{for } a < 2 \quad (0-1)$$

Here $2 = \lim_{N \rightarrow \infty} \mathbb{E}(\lambda_{\max})$. It is proved, that $\mathcal{F}_N(a)$ has an expansion in $1/N^2$, either in [31] by probabilistic methods, or in [16] by Riemann-Hilbert asymptotic analysis for orthogonal polynomials [5, 13, 14, 15]:

$$\mathcal{F}_N(a) = A_N + \sum_{g \geq 0} N^{2-2g} F^g(a) \quad (0-2)$$

with $F^g(a)$ independent of N . It is known that the $F^g(a)$ are the symplectic invariants introduced in [18] (see Section 1.1 below). To be short, they can be computed recursively, purely by means of algebraic geometry on a certain plane curve $\Sigma(a)$.

In [7], we have computed those coefficients $F^g(a)$, also for the generalization to β ensembles ($\beta > 0$) with polynomial potential. In this article, we shall consider only the hermitian case ($\beta = 2$). Because of universality, we may restrict ourselves to a quadratic potential, i.e. to the GUE ensemble.

Then, heuristically, one can study not so large deviations by plugging $a = 2 + N^{-2/3}s$ ($s < 0$) in Eqn. 0-2. The result is that (see [7]), with this scaling, each term of the expansion becomes of order 1:

$$\mathcal{F}_N(a = 2 + N^{-2/3}s) \sim A_{\text{GUE}} + \sum_{g \geq 0} (-s/2)^{3(1-g)} F^g(\Sigma_{\text{TW}}) \quad (0-3)$$

A_{GUE} is a constant that one can compute, and $F^g(\Sigma_{\text{TW}})$ are the symplectic invariants of a certain plane curve Σ_{TW} , which is a limit of the curve $\Sigma(a)$ when $a \rightarrow 2$. This limit object is in fact universal (it is the same if we start from a generic potential), it is the curve of equation $y^2 = x + \frac{1}{x} - 2$, up to normalization of x and y . Heuristically, it is expected that the RHS of Eqn. 0-3 is the exact $s \rightarrow -\infty$ asymptotic of $\lim_{N \rightarrow \infty} \mathcal{F}_N(a = 2 + N^{-2/3}s)$. Indeed, this method for $\beta > 0$ reproduces all previously known results [11, 30, 12, 2] on the left tail asymptotics of β Tracy-Widom laws. It also predicts the constant for any $\beta > 0$, and provides an algorithm to compute the expansion at all orders. In our case, for GUE:

$$\mathcal{P}_{\text{GUE}}^*(s) = 2^{1/24} e^{\zeta'(-1)} \exp \left(-\frac{|s|^3}{12} - \frac{\ln |s|}{8} + \sum_{g \geq 2} (-s/2)^{3(1-g)} F^g(\Sigma_{\text{TW}}) \right) \quad (0-4)$$

From Tracy and Widom [32], we know that $\mathcal{P}_{\text{GUE}}^*(s) = F_{\text{GUE}}(s)$, where F_{GUE} is related to Painlevé II equation, which itself appears in relation to an integrable system. However, by performing a $1/N$ expansion and computing F^g , the integrability structure is not manifest. At first, it was not clear why the asymptotic series in the RHS of Eqn. 0-4 should be related to Painlevé II. In this article, we fill this gap and give a proof of the following:

Proposition 0.1

$$\exp \left(-\frac{|s|^3}{12} - \frac{\ln |s|}{8} + \sum_{g \geq 2} (-s/2)^{3(1-g)} F^g(\Sigma_{\text{TW}}) \right) \quad (0-5)$$

is the asymptotic when $s \rightarrow -\infty$ of a Tau function for two 2×2 compatible systems of ODE:

$$\partial_x \Psi(x, s) = \mathbf{L}(x, s) \Psi(x, s), \quad \partial_s \Psi(x, s) = \mathbf{M}(x, s) \Psi(x, s) \quad (0-6)$$

whose compatibility condition is equivalent to a Painlevé II equation:

$$q''(s) = 2q^3(s) + sq(s) \quad (0-7)$$

The identity of Eqn. 0-3 was based on a heuristic argument, the double scaling limit of a matrix model, which would require a mathematical proof. In this article, we start from the RHS given by heuristics (i.e. from the world of the topological recursion), and we prove that it is indeed the expansion of the LHS. We do not use a Fredholm determinant representation of $\text{Prob}[\lambda_{\max} \leq a]$.

Outline

In the first part of the article, we gather older facts relevant for our problem:

- the framework of the topological recursion and its properties (Section 1.1-1.2).
- the general relation between 2×2 integrable systems and loop equations (Section 1.3-1.5).

In a second part, we refine the technical hypothesis underlying this very relation (the outcome is Theorem 2.1, our main theoretical result). We explain what really has to be checked (Section 2.1), and we sketch how this be done in practice (Section 2.3).

The third part is devoted to the application of these ideas to an integrable system related to Painlevé II (Sections 3.1-3.2). Subsequently, we prove Prop. 0.1 (Sections 3.3-3.5).

Let us mention that many ideas used in the proof also appeared in the work of M. Cafasso and O. Marchal [9] with different purposes (they related the $(2m, 1)$ minimal models appearing in the merging of two cuts in the 1 hermitian matrix model, to a hierarchy of equations containing Painlevé II), and in fact have their origin in the works of the second author with M. Bergère [3, 4]. In this article, our goal is to close (in the hermitian case only) the alternative approach to Tracy-Widom laws presented in [7].

Acknowledgments

We would like to thank M. Bergère for sharing his computations on Schrödinger equations and frequent valuable discussions, T. Grava for discussions about PII, O. Marchal and A. Its for comments, S.N. Majumdar, C. Nadal with whom we collaborated on a previous related work, and G. Schehr for motivating discussions. G.B. thanks the hospitality of the SISSA and of the Department of Maths and Statistics of Melbourne University, where part of this work was completed, and the organizers of the MSRI semester on Random Matrix Theory. The work of B.E. is partly supported by the ANR project GranMa "Grandes Matrices Aléatoires" ANR-08-BLAN-0311-01, by the European Science Foundation through the Misgam program, by the Québec government with the FQRNT. He also would like to thank the CRM (Centre de recherches mathématiques de Montréal, QC, Canada) and the CERN for its hospitality.

1 Basics around the topological recursion

1.1 Definitions

The topological recursion and symplectic invariants were axiomatically defined in [18], and we refer to [19] for a review, and to Appendix B for a summarized definition. It consists in an algorithm associating some numbers $F^g(\Sigma)$ and differential forms $\omega_n^g(\Sigma)(z_1, \dots, z_n)$ to a regular spectral curve Σ .

For our purposes, we call *spectral curve* the data of:

- a plane curve (\mathcal{C}, x, y) , in other words a Riemann surface \mathcal{C} , with two meromorphic functions x and y , $\mathcal{C} \mapsto \mathbb{C}$.
- a maximal open domain $U \subseteq \mathcal{C}$ on which x is a coordinate patch, called *physical sheet*.
- a Bergman kernel $B(z_1, z_2)$, i.e. a differential form in $z_1 \in \mathcal{C}$ and in $z_2 \in \mathcal{C}$, such that, in any local coordinate ξ :

$$B(z_1, z_2) \underset{z_1 \rightarrow z_2}{=} \frac{d\xi(z_1)d\xi(z_2)}{(\xi(z_1) - \xi(z_2))^2} + O(1) \quad (1-1)$$

The zeroes of dx are called *branchpoints* (name them a_i), and a spectral curve is said to be *regular* when these zeroes are simple and are not zeroes of dy . In other words, when $\sqrt{x - x(a_i)}$ is a good coordinate around $a_i \in \mathcal{C}$, and when y behaves like $y(z) \sim y(a_i) + y'(a_i) \sqrt{x(z) - x(a_i)}$ with $y'(a_i) \neq 0$.

We shall not give the full definitions of $F^g(\Sigma) = \omega_0^g(\Sigma)$ and $\omega_n^g(\Sigma)$ here, and rather refer to Appendix B or [18]. Their construction is axiomatic and relies only on algebraic geometry on \mathcal{C} . The construction is made by recursion on $2g - 2 + n$.

We rather mention the essential properties that we use here:

- $F^g(\Sigma)$ is invariant under any transformation $(x, y) \rightarrow (x_o, y_o)$ such that $|dx \wedge dy| = |dx_o \wedge dy_o|$. For this reason, the $F^g(\Sigma)$ are called *symplectic invariants*.
- For $2 - 2g - n < 0$, $\omega_n^g(\Sigma) \in T^*(\mathcal{C}) \otimes \dots \otimes T^*(\mathcal{C})$, i.e. $\omega_n^g(\Sigma)(z_1, \dots, z_n)$ is a meromorphic differential form in each $z_i \in \mathcal{C}$, symmetric in all z_i 's, and it has poles only at the branchpoints, of maximal degree $2(3g + n - 2)$, with vanishing residue.
- for $2 - 2g - n < 0$, $\omega_n^g(\Sigma)$ are homogeneous of degree $2 - 2g - n$, i.e. if we change $y \rightarrow \lambda y$, we have $\omega_n^g(\Sigma) \rightarrow \lambda^{2-2g-n} \omega_n^g(\Sigma)$, and $F^g(\Sigma) \rightarrow \lambda^{2-2g} F^g(\Sigma)$.

- $\omega_n^g(\Sigma)$ have nice scaling properties when the spectral curve approaches a singular spectral curve. We will be more precise when needed.
- $\omega_n^g(\Sigma)$ have nice properties under variation of the spectral curve. If Σ_t is a 1-parameter family of spectral curves, let us write $(\partial_t y dx - \partial_t x dy) = \Omega$, and represent the differential form Ω with the Bergman kernel:

$$\Omega(z) = \int_{\Omega^*} \Lambda_\Omega(\xi) B(\xi, z) \quad (1-2)$$

(this is form-cycle duality: Ω^* is the cycle dual to the differential form Ω , and in case the cycle is not regular, it is accompanied by a Jacobian Λ_Ω). Then:

$$\partial_t \omega_n^g(\Sigma_t)(z_1, \dots, z_n) = \int_{\Omega^*} \Lambda_\Omega(\xi) \omega_{n+1}^g(\Sigma_t)(\xi, z_1, \dots, z_n). \quad (1-3)$$

1.2 Matrix models and topological recursion

The topological recursion allows to find all order asymptotics of the large N expansion of matrix models. Let us recall how it works for the 1 hermitian matrix model with one hard edge, relevant for the application to Tracy-Widom law. For any potential V such that $|\int_{\mathbf{R}} d\xi e^{-\epsilon V(\xi)}| < \infty$ for some positive ϵ , let us consider N real random variables $\lambda_1, \dots, \lambda_N$ with joint probability law of density:

$$d\mu_N(\lambda) = \frac{1}{Z_N(\infty)} \prod_{1 \leq i < j \leq N} |\lambda_i - \lambda_j|^2 \cdot \prod_{i=1}^N d\lambda_i \exp(-NV(\lambda_i)) \quad (1-4)$$

The probability of having $\max \lambda_i \leq a$ is given by $\mathcal{P}(a) = Z_N([-\infty, a])/Z_N(\infty)$, where:

$$Z_N(J) = \int_{J^N} d\mu_N(\lambda) \quad (1-5)$$

$d\mu_N$ is the probability distribution induced on the eigenvalues of a random hermitian matrix of size $N \times N$, with probability law $dM \exp(-\frac{N}{t} \text{Tr } V(M))$ up to normalization. dM here is the canonical Lebesgue measure on the real vector spaces of hermitian matrices. The observables invariant under conjugation are spanned by the cumulants:

$$W_n(x_1, \dots, x_n) = \left\langle \prod_{i=n}^N \left(\sum_{j_i=1}^N \frac{1}{x_i - \lambda_{j_i}} \right) \right\rangle_c \quad (1-6)$$

When $\ln Z_N(a)$ admits an asymptotic expansion in powers of $1/N^2$, we write:

$$Z_N(a) = C_N \exp \left(\sum_{g \geq 0} N^{2-2g} F^g \right) \quad (1-7)$$

$$W_n(x_1, \dots, x_n) = \sum_{g \geq 0} N^{2-2g-n} W_n^g(x_1, \dots, x_n) \quad (1-8)$$

where C_N is a normalization constant depending only on N , and F^g and W_n^g are independent of N but depend on a and V . Before coming to the coefficients themselves, let us say that when the density of eigenvalues has a large N limit, $\rho(\xi)d\xi$, which admits a single interval $J_0 \subseteq]-\infty, a[$ as support, it is proved that such an expansion exists [31, 16]. If the support consists of several intervals, no such expansion exists (there are fast oscillatory terms depending on N), but there is still a way to extract the partition function from the topological recursion [20]. Yet, this discussion is out of the scope of this article. The theorem proved in [17, 10] is that when the large N expansion exists, then:

$$F^g = F^g(\Sigma(a)) \quad (1-9)$$

$$W_n^{g,k}(x(z_1), \dots, x(z_n))dx(z_1) \cdots dx(z_n) = \omega_n^{g,k}(\Sigma(a))(z_1, \dots, z_n) \quad (1-10)$$

for the spectral curve $\Sigma(a)$ determined by the equation $y = V'(x)/2 - W_1^0(x)$ and the Bergman kernel determined by:

$$B(z_1, z_2) = \left(W_2^0(x(z_1), x(z_2)) + \frac{1}{(x(z_1) - x(z_2))^2} \right) dx(z_1)dx(z_2) \quad (1-11)$$

$y(x)$ is analytic on the x complex plane, except on the eigenvalue support J_0 , where it admits a discontinuity $y(x + i0^+) - y(x - i0^-) = 2i\pi\rho(x)$. \mathcal{C} is a Riemann surface on which y can be analytically continued.

1.3 2×2 differential system and topological recursion

To any 2×2 first order differential system:

$$\partial_x \Psi = \mathbf{L} \Psi, \quad \Psi = \begin{pmatrix} \psi & \phi \\ \bar{\psi} & \bar{\phi} \end{pmatrix} \quad (1-12)$$

is associated the integrable kernel:

$$\mathcal{K}(x_1, x_2) = \frac{\psi(x_1)\bar{\phi}(x_2) - \bar{\psi}(x_1)\phi(x_2)}{x_1 - x_2} \quad (1-13)$$

We shall restrict ourselves to $\text{Tr } \mathbf{L} = 0$. So, $\partial_x(\det \Psi) = 0$, and we can choose the normalization:

$$\det \Psi = 1 \quad (1-14)$$

In [3] were also introduced correlators $\bar{\mathcal{W}}_n(x_1, \dots, x_n)$ and connected correlators $\mathcal{W}_n(x_1, \dots, x_n)$. The connected correlators are defined by:

$$\mathcal{W}_1(x) = \lim_{x' \rightarrow x} \left(\mathcal{K}(x, x') - \frac{1}{x - x'} \right) \quad (1-15)$$

$$\mathcal{W}_2(x_1, x_2) = -\mathcal{K}(x_1, x_2)\mathcal{K}(x_2, x_1) - \frac{1}{(x_1 - x_2)^2} \quad (1-16)$$

$$\mathcal{W}_n(x_1, \dots, x_n) = (-1)^{n+1} \sum_{\sigma \text{ cycles of } \mathfrak{S}_n} \prod_{i=1}^n \mathcal{K}(x_i, x_{\sigma(i)}) \quad (1-17)$$

and the correlators by:

$$\overline{\mathcal{W}}_n(x_1, \dots, x_n) = \text{"det"} \mathcal{K}(x_i, x_j) \quad (1-18)$$

where "det" means that each occurrence of $\mathcal{K}(x_i, x_i)$ in the determinant should be replaced by $\mathcal{W}_1(x_i)$, and each occurrence of $\mathcal{K}(x_i, x_j)\mathcal{K}(x_j, x_i)$ by $-\mathcal{W}_2(x_i, x_j)$. In other words the \mathcal{W}_n are the cumulants of the $\overline{\mathcal{W}}_n$.

For example:

$$\mathcal{W}_1 = \psi \partial_x \overline{\phi} - \overline{\psi} \partial_x \phi = -(\partial_x \psi \overline{\phi} - \partial_x \overline{\psi} \phi) \quad (1-19)$$

Eqn. 1-14 implies that all correlators are symmetric in the x_i 's. It can be checked that they do not have poles at coinciding points $x_i = x_j$, $i \neq j$. The spectral curve of a first order differential system is defined by the plane curve \mathcal{C} of equation:

$$\det(y - \mathbf{L}(x)) = 0. \quad (1-20)$$

Theorem 1.1 (proved in [3]) *Assume that:*

- (i) \mathbf{L} depends on some parameter N , and has a limit when $N \rightarrow \infty$.
- (ii) The spectral curve Σ_N of the system Eqn. 1-20 has a large N limit Σ_∞ which is regular, and has genus 0.
- (iii) $\mathcal{W}_n(x_1, \dots, x_n)$ admits an asymptotic expansion when $N \rightarrow \infty$ of the form $\mathcal{W}_n = \sum_{g \geq 0} N^{1-2g} \mathcal{W}_n^g$, and for $2g - 2 + n > 0$, $\mathcal{W}_n^g(x_1, \dots, x_n)$ may have singularities only at branchpoints of Σ_∞ .

Then, the expansion coefficients of the correlators are computed by the topological recursion applied to the spectral curve defined by Σ_∞ with Bergman kernel $B(z_1, z_2) = (\mathcal{W}_2^0(x(z_1), x(z_2)) + 1/(x(z_1) - x(z_2))^2) dx(z_1) dx(z_2)$:

$$\mathcal{W}_n^g(x(z_1), \dots, x(z_n)) dx(z_1) \cdots dx(z_n) = \omega_n^g(\Sigma_\infty)(z_1, \dots, z_n) \quad (1-21)$$

We stress that it is the large N limit of the spectral curve (also called "classical", or "semiclassical" spectral curve) which is relevant in this theorem. A similar result holds when Σ_∞ is not of genus 0 under an extra hypothesis, but this will not be needed here. So far, we did not speak of an integrable system, the theorem itself tells something

about one differential system alone. However, in many examples and in this article, proving that (iii) holds is made possible by the existence of other compatible systems $\partial_{t_j}\Psi = \mathbf{M}_j\Psi$, where t_j is a parameter of \mathbf{L} , and \mathbf{M}_j is rational in x . So, we shall see in Sections 2.1-2.4 that hypothesis (iii) can be considerably weakened.

1.4 Generalities on the Schrödinger equation

Let us write:

$$\mathbf{L} = \begin{pmatrix} a & b \\ c & -a \end{pmatrix} \quad (1-22)$$

Eqn. 1-12 imply that ψ and ϕ are two independent solutions of a second order equation:

$$\partial_x^2 \psi + B \partial_x \psi + V \psi = 0 \quad (1-23)$$

where:

$$B = -\frac{\partial_x b}{b} \quad (1-24)$$

$$V = \det \mathbf{L} - \partial_x a + a \frac{\partial_x b}{b} \quad (1-25)$$

All the same, $\bar{\psi}$ and $\bar{\phi}$ are independent solutions of the second order equation:

$$\partial_x^2 \bar{\psi} + \bar{B} \partial_x \bar{\psi} + \bar{V} \bar{\psi} = 0 \quad (1-26)$$

where:

$$\bar{B} = -\frac{\partial_x c}{c} \quad (1-27)$$

$$\bar{V} = \det \mathbf{L} + \partial_x a - a \frac{\partial_x c}{c} \quad (1-28)$$

We shall use a classical change of function in the theory of Schrödinger equations to study the asymptotics of Ψ at singularities of \mathbf{L} . We are grateful to M. Bergère who shared his experience on that subject with us. We can always write:

$$\psi = \text{cte} \sqrt{\frac{bf}{2}} \exp \left(- \int^x \frac{d\xi}{f(\xi)} \right) \quad (1-29)$$

$$\bar{\psi} = \overline{\text{cte}} \sqrt{\frac{c\bar{f}}{2}} \exp \left(- \int^x \frac{d\xi}{\bar{f}(\xi)} \right) \quad (1-30)$$

Then, Eqns. 1-23-1-26 are equivalent to:

$$\frac{1}{2} \partial_x^3 f = 2U(\partial_x f) + (\partial_x U)f \quad (1-31)$$

$$\frac{1}{2} \partial_x^3 \bar{f} = 2\bar{U}(\partial_x \bar{f}) + (\partial_x \bar{U})\bar{f} \quad (1-32)$$

with U, \bar{U} given by $-V, -\bar{V}$ up to a schwartzian derivative:

$$U = -V + \frac{3}{4} \frac{(\partial_x b)^2}{b^2} - \frac{1}{2} \frac{\partial_x^2 b}{b} \quad (1-33)$$

$$\bar{U} = -\bar{V} + \frac{3}{4} \frac{(\partial_x c)^2}{c^2} - \frac{1}{2} \frac{\partial_x^2 c}{c} \quad (1-34)$$

1.5 Tau function and topological recursion

Here, we modify slightly the systems by introducing a parameter $1/N$ in front of each derivative (very often, N turns out to be redundant with other parameters). For a large class of 2×2 first order compatible systems (for example this includes the $(p, 2)$ minimal models which are finite reductions of KdV) of the type:

$$\frac{1}{N} \partial_x \Psi = \mathbf{L} \Psi \quad \frac{1}{N} \partial_{t_j} \Psi = \mathbf{M}_j \Psi \quad (1-35)$$

and under the hypothesis of Thm. 1.1, it was stated in [4, 9] without details that $\exp\left(\sum_{g \geq 0} N^{2-2g} F^g(\Sigma_\infty)\right)$ is a formal Tau function. We shall make the argument explicit here. We are considering *any* system like Eqn. 1-35, such that \mathbf{L} and \mathbf{M}_j are rational fractions in x . We assume that \mathbf{L} has a large N limit $\mathbf{L}^{(0)}$, and an expansion in powers of $1/N$.

1.5.1 Definition of the τ -function

Here, we consider that N is a fixed parameter. To any solution $\Psi(x, \vec{t})$ of Eqn. 1-35, Jimbo-Miwa-Ueno [26] associate a Tau function $\tau(\vec{t})$ as follows. Let us consider the behavior of $\Psi(x, \vec{t})$ near each pole x_α of $\mathbf{L}(x, \vec{t})$, and write the $x \rightarrow x_\alpha$ asymptotics (valid in some angular sector near x_α):

$$\Psi(x, \vec{t}) = \tilde{\Psi}_\alpha(x, \vec{t}) \exp(\mathbf{T}_\alpha(x, \vec{t})) \quad (1-36)$$

where $e^{\mathbf{T}_\alpha}$ contains the essential singularity and the pole of $\Psi(x, \vec{t})$, and $\tilde{\Psi}_\alpha(x, \vec{t})$ is analytical when $x \rightarrow x_\alpha$. Then, the compatibility of the system Eqn. 1-35 implies that there exists a function $\tau(\vec{t})$ such that:

$$\partial_{t_j} \ln \tau = - \sum_{\alpha} \operatorname{Res}_{x \rightarrow x_\alpha} dx \operatorname{Tr} \tilde{\Psi}_\alpha^{-1} (\partial_x \tilde{\Psi}_\alpha) (\partial_{t_j} \mathbf{T}_\alpha) \quad (1-37)$$

$$= - \sum_{\alpha} \operatorname{Res}_{x \rightarrow x_\alpha} dx \operatorname{Tr} (\Psi^{-1} \partial_x \Psi) e^{-\mathbf{T}_\alpha} (\partial_{t_j} \mathbf{T}_\alpha) e^{\mathbf{T}_\alpha} \quad (1-38)$$

Proposition 1.1 *Assume that $\mathbf{T}_\alpha(x, \vec{t})$ is family of diagonal matrices, and write $\mathbf{T}_\alpha = T_\alpha \sigma_3 + c \mathbf{1}$ (where $\sigma_3 = \operatorname{diag}(1, -1)$). We have:*

$$\partial_{t_j} \ln \tau = 2 \sum_{\alpha} \operatorname{Res}_{x \rightarrow x_\alpha} dx (\partial_{t_j} T_\alpha) \mathcal{W}_1 \quad (1-39)$$

with the correlator $\mathcal{W}_1(x) = (\partial_x \psi) \bar{\phi} - (\partial_x \bar{\psi}) \phi$ introduced in Section 1.3.

proof:

$\Psi^{-1}\partial_x\Psi = \Psi^{-1}\mathbf{L}\Psi$ is traceless like \mathbf{L} , and $\mathbf{T}_\alpha(x, \vec{t})$ is a family of commuting matrices. Thus, only the component of \mathbf{T} of the Pauli matrix σ_3 is relevant. Eqn. 1-38 yields:

$$\partial_{t_j} \ln \tau = -2 \sum_{\alpha} \text{Res}_{x \rightarrow x_{\alpha}} dx (\partial_{t_j} T_{\alpha}) (\Psi^{-1} \partial_x \Psi)_{11} \quad (1-40)$$

And we compute using $\det \Psi = 1$:

$$\Psi^{-1} \partial_x \Psi = \begin{pmatrix} \bar{\phi} & -\phi \\ -\bar{\psi} & \psi \end{pmatrix} \cdot \begin{pmatrix} \partial_x \psi & \partial_x \phi \\ \partial_x \bar{\psi} & \partial_x \bar{\phi} \end{pmatrix} \quad (1-41)$$

$$= \begin{pmatrix} \partial_x \psi \bar{\phi} - \partial_x \bar{\psi} \phi & \bar{\phi} \partial_x \phi - \phi \partial_x \bar{\phi} \\ \psi (\partial_x \bar{\psi}) - \bar{\psi} \partial_x \psi & -\bar{\psi} \partial_x \phi + \psi \partial_x \bar{\phi} \end{pmatrix} \quad (1-42)$$

$$= \begin{pmatrix} -\mathcal{W}_1 & \bar{\phi} \partial_x \phi - \phi \partial_x \bar{\phi} \\ \psi \partial_x \bar{\psi} - \bar{\psi} \partial_x \psi & \mathcal{W}_1 \end{pmatrix} \quad (1-43)$$

□

1.5.2 Large N limit

We use Section 1.4 and keep the same notations, except that now, each derivative comes with a power of $1/N$. This modification can be easily traced back:

$$\psi(x) = \text{cte} \sqrt{\frac{b(x)f(x)}{2}} \exp \left(-N \int^x \frac{d\xi}{f(\xi)} \right) \quad (1-44)$$

$$\bar{\psi}(x) = \overline{\text{cte}} \sqrt{\frac{c(x)\bar{f}(x)}{2}} \exp \left(-N \int^x \frac{d\xi}{\bar{f}(\xi)} \right) \quad (1-45)$$

where f and \bar{f} satisfy:

$$\frac{1}{2N^2} \partial_x^3 f = 2U(\partial_x f) + (\partial_x U)f \quad (1-46)$$

$$\frac{1}{2N^2} \partial_x^3 \bar{f} = 2\bar{U}(\partial_x \bar{f}) + (\partial_x \bar{U})\bar{f} \quad (1-47)$$

with the potential:

$$U = -\det \mathbf{L} + \frac{1}{N} \left(\partial_x a - a \frac{\partial_x b}{b} \right) + \frac{1}{N^2} \left(\frac{3}{4} \frac{(\partial_x b)^2}{b^2} - \frac{1}{2} \frac{\partial_x^2 b}{b} \right) \quad (1-48)$$

$$\bar{U} = -\det \mathbf{L} + \frac{1}{N} \left(-\partial_x a + a \frac{\partial_x c}{c} \right) + \frac{1}{N^2} \left(\frac{3}{4} \frac{(\partial_x c)^2}{c^2} - \frac{1}{2} \frac{\partial_x^2 c}{c} \right) \quad (1-49)$$

Let us write with superscript ⁰ the large N limit of these quantities. In particular:

$$\mathbf{L}^{(0)} = \begin{pmatrix} a^{(0)} & b^{(0)} \\ c^{(0)} & -a^{(0)} \end{pmatrix} \quad (1-50)$$

Recall that we defined the spectral curve through the equation $\det(y - \mathbf{L}(x)) = 0$. Since \mathbf{L} is traceless, it is equivalent to $y^2 = -\det \mathbf{L}(x)$. Hence, the large N limit of the spectral curve has equation:

$$\Sigma_\infty(\vec{t}) : \quad y^2 = -\det \mathbf{L}^{(0)}(x) \quad (1-51)$$

Let us define $y(x)$ by this equation, with a choice of the branch of the square root depending on the asymptotic $x \rightarrow x_\alpha$ we require to define ψ (see Eqn. 1-54 below). According to Eqns. 1-48, the potential in the large N limit is given by:

$$U^{(0)}(x) = \overline{U}^{(0)}(x) = y^2(x) \quad (1-52)$$

And, from Eqn. 1-46, we get:

$$f^{(0)}(x) \propto 1/y(x), \quad \overline{f}^{(0)}(x) \propto 1/y(x) \quad (1-53)$$

Matching the asymptotics gives the proportionality constants and we get:

$$\psi^{(0)}(x) = \text{cte} \sqrt{\frac{b^{(0)}(x)}{2y(x)}} \exp \left(-N \int_o^x y(\xi) d\xi \right), \quad (1-54)$$

$$\overline{\phi}^{(0)}(x) = \overline{\text{cte}} \sqrt{\frac{c^{(0)}(x)}{2y(x)}} \exp \left(N \int_o^x y(\xi) d\xi \right). \quad (1-55)$$

1.5.3 $1/N$ expansion of the τ -function

We now want to study asymptotics when $N \rightarrow \infty$. We assume that

$$\mathbf{T}_\alpha(x) = N \mathbf{T}_\alpha^{(0)}(x) = -N \sigma_3 \left(\int_o^x y(\xi) d\xi \right)_{-, \alpha} \quad (1-56)$$

where $(\cdots)_{-, \alpha}$ is the divergent part when $x \rightarrow x_\alpha$, and o is an arbitrary origin of integration. In other words, we assume that $\mathbf{L}(x)$ and $\mathbf{L}^{(0)}(x)$ have the same poles $\{x_\alpha\}$, that the behavior of Ψ at these poles is entirely given by the large N limit of the system. This happens quite often in practice. Then, we rewrite Eqn. 1-39:

$$\partial_{t_j} \tau = -2N \sum_\alpha \text{Res}_{x \rightarrow x_\alpha} \left(\int_o^x \partial_{t_j} y(\xi) d\xi \right) \mathcal{W}_1(x) \quad (1-57)$$

Provided the hypothesis of Thm. 1.1 hold, the topological recursion can be applied, with the spectral curve $(\Sigma_\infty(\vec{t}))$ defined in Eqn. 1-51. Let us compare to $\mathcal{F}(\vec{t})$ defined by the large N asymptotic series:

$$\mathcal{F}(\vec{t}) \equiv \sum_{g \geq 0} N^{2-2g} F^g(\Sigma_\infty(\vec{t})) \quad (1-58)$$

We also define:

$$W_1(x) \equiv \sum_{g \geq 0} N^{1-2g} W_1^g(\Sigma_\infty(\vec{t}))(x) \quad (1-59)$$

Let us call \mathcal{C} , the underlying curve of $\Sigma_\infty(\vec{t})$, and B its Bergman kernel. We shall use the properties of the topological recursion (Section 1.1) to compute the variations of $\mathcal{F}(\vec{t})$ wrt t_j . We have to represent

$$(\partial_{t_j} y \, dx)(\xi) = \operatorname{Res}_{z \rightarrow \xi} B(z, \xi) \int_o^z y \, dx \quad (1-60)$$

Notice that the RHS do not depend on o . For 2×2 systems, $\Sigma_\infty(\vec{t})$ is hyperelliptic. So, each pole x_α of $\int_o^z y \, dx$ has two preimages z_α and \bar{z}_α in \mathcal{C} . One can move the contour to surround these points:

$$(\partial_{t_j} y \, dx)(\xi) = - \sum_{\alpha} \operatorname{Res}_{z \rightarrow z_\alpha, \bar{z}_\alpha} B(z, \xi) \int_o^z \partial_{t_j} y \, dx \quad (1-61)$$

Subsequently:

$$\partial_{t_j} F^g(\Sigma_\infty(\vec{t})) = - \sum_{\alpha} \operatorname{Res}_{z \rightarrow z_\alpha, \bar{z}_\alpha} dx(z) W_1^g(z) \int_o^z \partial_{t_j} y \, dx \quad (1-62)$$

In the case of an hyperelliptic curve, the involution is a symmetry in the construction of W_1^g , and one finds that the residues at z_α and \bar{z}_α are equal. Hence:

$$\begin{aligned} \partial_{t_j} \mathcal{F}(\vec{t}) &= \sum_{g \geq 0} N^{2-2g} \partial_{t_j} F^g(\Sigma_\infty(\vec{t})) \\ &= - \sum_{\alpha} \operatorname{Res}_{z \rightarrow z_\alpha} dx(z) 2N W_1(x(z)) \int_o^z \partial_{t_j} y \, dx \\ &= \partial_{t_j} \ln \tau \end{aligned} \quad (1-63)$$

This proves the identification of the resummation of symplectic invariants with the τ function.

2 Refinement on the assumptions

2.1 A remark on hypothesis (iii)

Starting from any solution to $\frac{1}{N} \partial_x \Psi = \mathbf{L} \Psi$, with \mathbf{L} a given 2×2 matrix, rational in x and traceless, the connected correlators \mathcal{W}_n were defined in [3] and Section 1.3 such that they satisfy "loop equations". When these loop equations are written in terms of the τ function, they appear identical to Virasoro constraints³. Loop equations

³Independently of [3], Virasoro constraints have also been found in [27] in the case of 2×2 Schlesinger system.

can have many solutions, but their solution is unique if one requires some analyticity properties. In fact, provided there is a large parameter N in the problem, and provided an expansion:

$$\mathcal{W}_n = \sum_{g \geq 0} N^{2-2g-n} \mathcal{W}_n^g \quad (2-1)$$

exists, the topological recursion gives the unique solution of the hierarchy of loop equations satisfied by \mathcal{W}_n^g , whose analytical properties are specified by the data of the spectral curve.

The matter is then, for the Ψ one is interested in, to find the appropriate spectral curve Σ_∞ . It is often easy to prove that Ψ has an expansion in $1/N$, and to discuss its analytical properties (see Section 2.3). However, hypothesis (iii) of Thm. 1.1 is stronger. Indeed, the two essential properties of Eqn. 2-1 are:

- Fixed parity: only powers of $1/N$ with parity $(-1)^n$ appears in the expansion of \mathcal{W}_n .
- Factorization property: the leading order of \mathcal{W}_n is $O(N^{2-n})$.

We believe that in practice, "fixing parity" is only a matter of rescaling correctly with N the parameters of the differential system⁴ $\partial_x \Psi = \mathbf{L} \Psi$ and that $\mathbf{L}^{(0)}$. In Appendix A, we give an example based on a Lax pair associated to Painlevé II $_\alpha$ equation, where the fixed parity property does not hold if one does not rescale α as well.

Now, we argue that the factorization property is automatic, as soon as one realizes $\partial_x \Psi = \mathbf{L} \Psi$ as the member of a full integrable hierarchy. We present this argument for 2×2 systems with 1 pole at $x = \infty$ (the case of interest in this article), but most of these ideas could be adapted without difficulty to several poles and larger systems. We are currently working out the generalization of all arguments presented in this article to the case of $d \times d$ differential systems [8].

2.2 Loop insertion operator and properties

In some sector near $x = \infty$, we consider a solution Ψ whose asymptotic is given by $\Psi = \tilde{\Psi}_\alpha e^{N T \sigma_3}$, with $\tilde{\Psi}$ regular at $x = \infty$, and:

$$T = \sum_{j \geq 1} t_j x^j \quad (2-2)$$

for some constants t_j . It is a standard result in integrable systems [1] that t_j can be used to define commuting flows. The result yields a matrix \mathbf{L} and solution Ψ , which

⁴This is not a void statement, because Theorem 1.1 asks by convention for a $\frac{1}{N}$ in front of each ∂_x in the differential system, and also that $\lim_{N \rightarrow \infty} \mathbf{L}$ exists.

now depends on all t_j , with sectorwise asymptotic given by Eqn. 2-2, and such that:

$$\begin{aligned}\frac{1}{N} \partial_x \Psi &= \mathbf{L} \Psi \\ \frac{1}{N} \partial_{t_j} \Psi &= \mathbf{M}_j \Psi \\ \mathbf{M}_j(x) &= \left[\tilde{\Psi}(x) x^j \sigma_3 \tilde{\Psi}^{-1}(x) \right]_- \end{aligned} \quad (2-3)$$

$[\cdots]_{-,x}$ denotes the divergent part when $x \rightarrow \infty$, and obviously \mathbf{M}_j is a polynomial in x of degree j . Let us define the *loop insertion operator* as a Laurent series of operators when $x \rightarrow \infty$:

$$\delta_x = \sum_{j \geq 1} \frac{\partial_{t_j}}{x^{j-1}} \quad (2-4)$$

It is merely a way to collect all the flows⁵.

Lemma 2.1 *We write $\widetilde{\cdots}$ the functions ψ, ϕ, \cdots with their essential singularity removed. The following 2×2 matrix:*

$$\begin{aligned} \mathbf{P}(x) &\equiv \frac{\mathbf{1} - \tilde{\Psi}(x) \sigma_3 \tilde{\Psi}^{-1}(x)}{2} \\ &= \begin{pmatrix} -\bar{\psi}(x)\phi(x) & \tilde{\psi}(x)\tilde{\phi}(x) \\ -\tilde{\bar{\psi}}(x)\tilde{\phi}(x) & \psi(x)\bar{\phi}(x) \end{pmatrix} \end{aligned} \quad (2-5)$$

is a projector, and we have:

$$\frac{1}{N} \delta_{x_2} \tilde{\Psi}(x_1) = \frac{2(\mathbf{P}(x_2) - \mathbf{P}(x_1))}{x_2 - x_1} \tilde{\Psi}(x_1) \quad (2-6)$$

$$\frac{1}{N} \delta_{x_2} \mathbf{P}(x_1) = \frac{[\mathbf{P}(x_1), \mathbf{P}(x_2)]}{x_2 - x_1} \quad (2-7)$$

proof:

Eqn. 2-5 is a straightforward computation. We just notice, according to the representations of Eqns. 1-29-1-30, that the essential singularities cancels when we compute the cross-products: $\psi \bar{\phi} = \tilde{\psi} \tilde{\bar{\phi}}$ and $\bar{\psi} \phi = \tilde{\bar{\psi}} \tilde{\phi}$. Then, we observe that \mathbf{P} is a rank 1 matrix:

$$\mathbf{P} = {}^t(\tilde{\phi} \quad \tilde{\bar{\phi}}) (\tilde{\psi} \quad \tilde{\bar{\psi}}) \mathbf{J} \quad (2-8)$$

with:

$$\mathbf{J} = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}, \quad \mathbf{J}^2 = -1, \quad {}^t \mathbf{J} = \mathbf{J}^{-1} = -\mathbf{J} \quad (2-9)$$

By definition $\text{Tr } \mathbf{P} = 1$. Hence, $\mathbf{P}^2 = (\text{Tr } \mathbf{P}) \mathbf{P} = \mathbf{P}$. Now, let us compute the action of the loop insertion operator on Ψ . $\partial_{t_j} \Psi = \mathbf{M}_j \Psi$ translates into:

$$\frac{1}{N} \partial_{t_j} \tilde{\Psi}(x_1) = \left[\tilde{\Psi}(x_1) x_1^j \sigma_3 \tilde{\Psi}^{-1}(x_1) \right]_- \tilde{\Psi}(x_1) - \tilde{\Psi}(x_1) x_1^j \sigma_3 \quad (2-10)$$

⁵ $\frac{1}{N} \delta_x$ given by Eqn. 2-4 gives a realization of the formal loop insertion operator introduced in [3]

It is implied that $[\cdots]_-$ is the divergent part when $x_1 \rightarrow \infty$. Doing the summation over j :

$$\begin{aligned}
\frac{1}{N} \delta_{x_2} \tilde{\Psi}(x_1) &= \left[\tilde{\Psi}(x_1) \frac{\sigma_3}{x_2 - x_1} \tilde{\Psi}^{-1}(x_1) \right]_- \tilde{\Psi}(x_1) - \tilde{\Psi}(x_1) \frac{\sigma_3}{x_2 - x_1} \\
&= \frac{\tilde{\Psi}(x_1) \sigma_3 \tilde{\Psi}^{-1}(x_1) - \tilde{\Psi}(x_2) \sigma_3 \tilde{\Psi}^{-1}(x_2)}{x_2 - x_1} \tilde{\Psi}(x_1) \\
&= \frac{2(\mathbf{P}(x_2) - \mathbf{P}(x_1))}{x_2 - x_1} \tilde{\Psi}(x_1)
\end{aligned}$$

We apply this result to compute:

$$\begin{aligned}
\frac{1}{N} \delta_{x_2} \mathbf{P}(x_1) &= -\frac{1}{2} \delta_{x_2} \left(\tilde{\Psi}(x_1) \sigma_3 \tilde{\Psi}(x_1)^{-1} \right) \\
&= -\frac{\mathbf{P}(x_2) - \mathbf{P}(x_1)}{x_2 - x_1} \tilde{\Psi}(x_1) \sigma_3 \tilde{\Psi}^{-1}(x_1) + \tilde{\Psi}(x_1) \sigma_3 \tilde{\Psi}^{-1}(x_1) \frac{\mathbf{P}(x_2) - \mathbf{P}(x_1)}{x_2 - x_1} \\
&= \frac{-(\mathbf{P}(x_2) - \mathbf{P}(x_1))\mathbf{P}(x_1) + \mathbf{P}(x_1)(\mathbf{P}(x_2) - \mathbf{P}(x_1))}{x_2 - x_1} \\
&= \frac{[\mathbf{P}(x_1), \mathbf{P}(x_2)]}{x_2 - x_1}
\end{aligned}$$

□

Corollary 2.1 *The integrable kernel is self-replicating:*

$$\frac{1}{N} \delta_{x_2} \mathcal{K}(x_1, x_3) = -\mathcal{K}(x_1, x_2) \mathcal{K}(x_2, x_3) \quad (2-11)$$

Proposition 2.1 *The correlators can be expressed in terms of the projector \mathbf{P} :*

$$\mathcal{W}_1(x) = -N \operatorname{Tr} \mathbf{P}(x) \mathbf{L}(x) \quad (2-12)$$

$$\mathcal{W}_2(x_1, x_2) = -\frac{1}{2} \frac{\operatorname{Tr} (\mathbf{P}(x_1) - \mathbf{P}(x_2))^2}{(x_1 - x_2)^2} \quad (2-13)$$

and for $n \geq 3$:

$$\mathcal{W}_n(x_1, \dots, x_n) = N^{2-n} (-1)^{n+1} \sum_{\sigma \text{ cycles of } \mathfrak{S}_n} \frac{\operatorname{Tr} \mathbf{P}(x_1) \mathbf{P}(x_{\sigma(1)}) \cdots \mathbf{P}(x_{\sigma^{n-1}(1)})}{(x_1 - x_{\sigma(1)})(x_{\sigma(1)} - x_{\sigma^2(1)}) \cdots (x_{\sigma^{n-1}(1)} - x_1)} \quad (2-14)$$

Or concisely with the loop insertion operator:

$$\mathcal{W}_n(x_1, \dots, x_n) = N^{-n} \delta_{x_1} \cdots \delta_{x_n} \ln \tau \quad (2-15)$$

As a consequence of the existence of a loop insertion operator and Eqn. 2-15, the factorization property is automatic, and the requirements of hypothesis (iii) have to be checked only on $\mathcal{W}_1(x)$.

2.3 Analyticity properties of the $1/N$ expansion

2.3.1 Summary of the problem

Let us start with two 2×2 matrices $\mathbf{L}(x, t)$ and $\mathbf{M}(x, t)$ such that:

- \mathbf{L} and \mathbf{M} are traceless, and rational in x .
- They are solution of the compatibility equation:

$$\frac{1}{N} \partial_t \mathbf{L} - \frac{1}{N} \partial_x \mathbf{M} + [\mathbf{L}, \mathbf{M}] = 0 \quad (2-16)$$

- They admit an asymptotic expansion when $N \rightarrow \infty$ of the form:

$$\mathbf{L} = \sum_{l \geq 0} N^{-l} \mathbf{L}^{(l)}, \quad \mathbf{M} = \sum_{l \geq 0} N^{-l} \mathbf{M}^{(l)} \quad (2-17)$$

and consider the equations:

$$\frac{1}{N} \partial_x \Psi = \mathbf{L} \Psi, \quad \frac{1}{N} \partial_t \Psi = \mathbf{M} \Psi \quad (2-18)$$

In other words, we first restrict ourselves to a hierarchy with only one time t . We shall discuss the analytic properties of ψ (the discussion would be similar for $\bar{\psi}$, ϕ and $\bar{\phi}$). From the differential system wrt x , we know (Section 1.5.2) that it admits a large N asymptotic expansion of the form:

$$\psi(x) = \left(\sum_{l \geq 0} N^{-l} \tilde{\psi}^{(l)}(x) \right) e^{-N \int_o^x y(\xi) d\xi} \quad (2-19)$$

where o is some integration constant which may depend on t . The leading order is given by:

$$\tilde{\psi}^{(0)} = \sqrt{\frac{\mathbf{L}_{21}}{2y}} \quad (2-20)$$

and y is determined by the large N spectral curve associated to the differential system in x :

$$\Sigma_\infty : \quad y^2 = -\det \mathbf{L}^{(0)} \quad (2-21)$$

Similarly, we may define the large N spectral curve associated to the differential system in t :

$$\underline{\Sigma}_\infty ; \quad \underline{y}^2 = -\det \mathbf{M}^{(0)} \quad (2-22)$$

We shall see that Σ_∞ and $\underline{\Sigma}_\infty$ play an important role to locate the singularities of ψ .

2.3.2 Analytical properties

Let us describe the analytical properties of $\tilde{\psi}^{(0)}$ in the variable x :

- It has branchpoints (order $\frac{-(2r+1)}{4}$, $r \in \mathbb{Z}$) at branchpoints of Σ_∞ .
- It has branchpoints (order $\frac{(2r+1)}{2}$, $r \in \mathbb{Z}$) at the zeroes or poles of $\frac{\mathbf{L}_{21}^{(0)}}{y}$ which are not branchpoints of Σ_∞ .

We can insert Eqn. 3-41 in one or the other differential system to determine recursively the subleading orders. It is more convenient to write the two second order equations satisfied by ψ :

$$\begin{aligned}\frac{1}{N^2} \partial_x^2 \psi + \frac{1}{N} B \partial_x \psi + V \psi &= 0 \\ \frac{1}{N^2} \partial_t^2 \psi + \frac{1}{N} \underline{B} \partial_t \psi + \underline{V} \psi &= 0\end{aligned}$$

where:

$$\begin{aligned}B &= -\frac{\partial_x b}{b} = \sum_{l \geq 0} N^{-l} B^{(l)} \\ V &= \det \mathbf{L} - \frac{1}{N} \partial_x \mathbf{L}_{11} + \frac{1}{N} \mathbf{L}_{11} \frac{\partial_x \mathbf{L}_{21}}{\mathbf{L}_{21}} = -y^2 + \sum_{l \geq 1} N^{-l} V^{(l)} \\ \underline{B} &= -\frac{\partial_x \mathbf{M}_{21}}{\mathbf{M}_{21}} = \sum_{l \geq 0} N^{-l} \underline{B}^{(l)} \\ \underline{V} &= \det \mathbf{M} - \frac{1}{N} \partial_t \mathbf{M}_{11} + \frac{1}{N} \mathbf{M}_{11} \frac{\partial_t \mathbf{M}_{21}}{\mathbf{M}_{21}} = -\underline{y}^2 + \sum_{l \geq 1} N^{-l} \underline{V}^{(l)}\end{aligned}\quad (2-23)$$

Knowing $\tilde{\psi}^{(0)}$, we can compute recursively, if we assume that $B^{(0)}, \underline{B}^{(0)}$ are not identically zero:

$$\begin{aligned}\tilde{\psi}^{(l+1)} &= \frac{1}{y B^{(0)}} \left[-\partial_x^2 \tilde{\psi}^{(l)} + (\partial_x y) \tilde{\psi}^{(l)} + 2y (\partial_x \tilde{\psi}^{(l)}) \right. \\ &\quad \left. \sum_{m=0}^l B^{(l-m)} \partial_x \tilde{\psi}^{(m)} - (V^{(l+1-m)} + y B^{(l+1-m)}) \tilde{\psi}^{(m)} \right]\end{aligned}\quad (2-24)$$

$$\begin{aligned}&= \frac{1}{y \underline{B}^{(0)}} \left[-\partial_t^2 \tilde{\psi}^{(l)} + (\partial_t y) \tilde{\psi}^{(l)} + 2y (\partial_t \tilde{\psi}^{(l)}) \right. \\ &\quad \left. + \sum_{m=0}^l \underline{B}^{(l-m)} \partial_t \tilde{\psi}^{(m)} - (\underline{V}^{(l+1-m)} + y \underline{B}^{(l+1-m)}) \tilde{\psi}^{(m)} \right]\end{aligned}\quad (2-25)$$

A priori, $\tilde{\psi}^{(l+1)}$ has singularities at points where $\tilde{\psi}^{(m)}$ (for $0 \leq m \leq l$) had already singularities, and at poles of \mathbf{L} and \mathbf{M} . Besides, if we consider the differential system wrt x , new singularities may appear at each step:

- At the branchpoints of (Σ_∞) (through $1/y$).
- At the zeroes (in the x variable) of $B^{(0)}$.

If we rather consider the differential system wrt t , new singularities may appear only:

- At the branchpoints of $\underline{\Sigma}_\infty$ (through $1/\underline{y}$).
- At the zeroes (in the x variable) of $\underline{B}^{(0)}$.

Subsequently, $\psi^{(l)}(x)$ may have singularities only:

- At the poles in the expansion of \mathbf{L} and \mathbf{M} .
- At the branchpoints of Σ_∞ .
- At the branchpoints of $\underline{\Sigma}_\infty$.
- At the common zeroes (in the x variable) of $B^{(0)}$ and $\underline{B}^{(0)}$.

In many examples (and in this article), the fourth point is the key ingredient to rule out singularities at the zeroes of y which are not branchpoints. It is often enough to have one time t , but the same argument could be repeated in presence of other times t_j , giving a priori more restrictions on the position of the singularities. Eventually, the singularities of \mathcal{W}_n can be inferred from their definition (Eqn. 1-17) and the singularities of $\tilde{\psi}^{(l)}$. We just notice that this definition does not introduce poles at coinciding points.

2.4 Conclusion

We may formulate a stronger version of Theorem 1.1.

Theorem 2.1 *Assume that:*

- (i) \mathbf{L} depends on some parameter N , and has a limit when $N \rightarrow \infty$.
- (ii) The spectral curve Σ_N of the system Eqn. 1-12 has a large N limit Σ_∞ which is regular, and has genus 0.
- (iii)' $\mathcal{W}_1(x)$ admits an asymptotic expansion when $N \rightarrow \infty$ of the form : $\mathcal{W}_1 = \sum_{g \geq 0} N^{1-2g} \mathcal{W}_1^g$, and $\mathcal{W}_1^g(x)$ for $g \neq 0$ may have singularities only at branchpoints of Σ_∞ .

Then, $\mathcal{W}_n(x_1, \dots, x_n)$ admits an expansion of the form:

$$\mathcal{W}_n = \sum_{g \geq 0} N^{2-2g-n} \mathcal{W}_n^g \quad (2-26)$$

The expansion coefficients of the correlators have only singularities at the branchpoints of Σ_∞ (for $2g - 2 + n > 0$), and are computed by the topological recursion applied to Σ_∞ :

$$\mathcal{W}_n^g(x(z_1), \dots, x(z_n)) dx(z_1) \cdots dx(z_n) = \omega_n^g(\Sigma_\infty)(z_1, \dots, z_n) \quad (2-27)$$

In practice, hypothesis (iii)' requires to check that:

- \mathcal{W}_1 has an expansion in odd powers of N . We think that it can always be achieved in an appropriate rescaling of the parameters of \mathbf{L} with N .
- \mathcal{W}_1^g has only singularities at branchpoints of Σ_∞ . This is not too difficult, because we know where singularities can arise in Ψ (Section 2.3.2).

A similar result holds when Σ_∞ is not of genus 0 under an extra hypothesis on \mathcal{W}_1^g , but is beyond of the scope of the present article.

3 Application to the Tracy-Widom GUE law

3.1 Tracy-Widom GUE and integrability

The Tracy-Widom GUE law $F_{\text{GUE}}(s)$ can be defined as a Fredholm determinant:

$$F_{\text{GUE}}(s) = \det(\mathbf{1} - \mathbf{K}_{\text{Ai}})_{L^2([s, \infty[)} \quad (3-1)$$

where \mathbf{K}_{Ai} is the Airy kernel:

$$(\mathbf{K}_{\text{Ai}} \cdot f)(x) = \int_{\mathbf{R}} dy \frac{\text{Ai}(x)\text{Ai}'(y) - \text{Ai}'(x)\text{Ai}(y)}{x - y} f(y) \quad (3-2)$$

In their celebrated article [32], Tracy and Widom have proved an alternative formula making the link with an integrable system:

$$F_{\text{GUE}}(s) = \exp \left(\int_s^\infty H(t) dt \right) \quad (3-3)$$

where $H(s) = -q^2(s)$, and $q(s)$ is the unique solution of the Painlevé II equation:

$$q''(s) = 2q^3(s) + sq(s) \quad (3-4)$$

which satisfies [25]:

$$q(s) \underset{s \rightarrow +\infty}{\sim} \text{Ai}(s) \sim \frac{\exp(-\frac{2}{3}s^{3/2})}{2\sqrt{\pi}s^{1/4}} \quad (3-5)$$

$H(s)$ can be identified with a Hamiltonian for PII [29], and $F_{\text{GUE}}(s)$ with a τ -function associated to this family of Hamiltonians ([24] in the sense of Okamoto, [6] in the sense of Jimbo-Miwa-Ueno).

The Painlevé II equation appears [22] as the compatibility condition of the following system for $\Psi(x, s)$:

$$\begin{cases} \partial_x \Psi &= \mathbf{L} \Psi \\ \partial_s \Psi &= \mathbf{M} \Psi \\ \Psi &= \tilde{\Psi} \exp \left[i \left(\frac{4}{3} x^3 + sx \right) \sigma_3 \right] \quad \text{when } x \rightarrow +\infty \end{cases} \quad (3-6)$$

where $\tilde{\Psi} = \mathbf{1} + O(1/x)$ when $x \rightarrow +\infty$. The Lax pair is given (\mathbf{L}, \mathbf{M}) is given by:

$$\begin{aligned} \mathbf{L}(x, s) &= \begin{pmatrix} -4ix^2 - i(s + 2q^2(s)) & 4xq(s) + 2ip(s) \\ 4xq(s) - 2ip(s) & 4ix^2 + i(s + 2q^2(s)) \end{pmatrix} \\ \mathbf{M}(x, s) &= \begin{pmatrix} -ix & q(s) \\ q(s) & ix \end{pmatrix} \end{aligned} \quad (3-7)$$

The necessary condition of existence of Ψ is $\partial_s \mathbf{L} - \partial_x \mathbf{M} + [\mathbf{L}, \mathbf{M}] = 0$. This implies that $q(s)$ is solution of PII, $p(s) = q'(s)$. The asymptotic behavior of Ψ determines the asymptotic behavior of Eqn. 3-6 for $q(s)$, picking up the Hastings-McLeod solution of PII [25]. The existence of the solution $\Psi(x, s)$ was shown in [22]. This system is an example of isomonodromy problem, itself closely related to a Riemann-Hilbert problem. Many authors have contributed to the study of these systems, and we refer to [23] for a review of the theory.

3.2 $s \rightarrow -\infty$ asymptotics and spectral curve

Let us introduce a redundant parameter N . We define:

$$\begin{cases} x = N^{1/3} X \\ s = N^{2/3} S \end{cases}, \quad q(s) = N^{1/3} Q(S) \quad (3-8)$$

Then, Eqn. 3-6 is equivalent to:

$$\begin{cases} \frac{1}{N} \partial_X \Psi &= \mathbf{L} \Psi \\ \frac{1}{N} \partial_S \Psi &= \mathbf{M} \Psi \\ \Psi &= \tilde{\Psi} \exp \left[iN \left(\frac{4}{3} X^3 + SX \right) \sigma_3 \right] \quad \text{when } X \rightarrow +\infty \end{cases} \quad (3-9)$$

with the Lax pair:

$$\begin{aligned} \mathbf{L}(X, S) &= \begin{pmatrix} -4iX^2 - i(S + 2Q^2(S)) & 4XQ(S) + \frac{2iQ'(S)}{N} \\ 4XQ(S) - \frac{2iQ'(S)}{N} & 4iX^2 + i(S + 2Q^2(S)) \end{pmatrix} \\ \mathbf{M}(X, S) &= \begin{pmatrix} -iX & Q(S) \\ Q(S) & iX \end{pmatrix} \end{aligned} \quad (3-10)$$

and the compatibility equation:

$$\frac{1}{N^2}Q''(S) = 2Q(S)^3 + SQ(S) \quad (3-11)$$

We are now in the situation described in Section 1, where each derivative appears with a prefactor $1/N$. Again, $Q(S)$ is given by the Hastings-McLeod solution to Painlevé II, which is the unique solution [25] such that, for $S < 0$, $\lim_{N \rightarrow \infty} Q(S)^2 = -\frac{S}{2}$. Because of this property, we may study the $s \rightarrow -\infty$ asymptotics for the system Eqn. 3-6 by studying the $N \rightarrow \infty$ asymptotics of the system of Eqn. 3-9. Then, the basic remark is that the derivative terms are subleading. After Eqn. 1-20, the finite N spectral curve for this system is:

$$Y^2 = -16X^4 - 8X^2S - S^2 - 4Q^4(S) - 4SQ^2(S) + \frac{4(Q'(S))^2}{N^2} \quad (3-12)$$

In the large N limit, since $Q^2(S) \sim -S/2$ (we assume $S < 0$), we obtain:

$$(\Sigma_\infty); \quad Y^2 = -16X^2 \left(X^2 + \frac{S}{2} \right) \quad (3-13)$$

It can be brought in a canonical form by rescaling:

$$(\widehat{\Sigma}) : \quad \widehat{y}^2 = \frac{1}{4}\widehat{x}^2(\widehat{x}^2 + 4) \quad (3-14)$$

where

$$\widehat{x} = i\sqrt{\frac{8}{-S}}X, \quad \widehat{y} = \frac{i}{S}Y \quad (3-15)$$

For this transformation, $YdX = (-S/2)^{3/2}\widehat{y}d\widehat{x}$. Going from the topological recursion of Σ_∞ to that of $\widehat{\Sigma}$ is only a matter of rescaling:

$$F^g(\Sigma_\infty) = (-S/2)^{3(1-g)} F^g(\widehat{\Sigma}) \quad (3-16)$$

$$\omega_n^{(g)}(\Sigma_\infty) = (-S/2)^{3(1-g-n/2)} \omega_n^g(\widehat{\Sigma}) \quad (3-17)$$

Thus:

$$\begin{aligned} \mathcal{F}(\Sigma_\infty) &= \sum_{g \geq 0} N^{2-2g} F^g(\Sigma_\infty) \\ &= \sum_{g \geq 0} (N^{2/3} (-S/2))^{3(1-g)} F^g(\widehat{\Sigma}) \\ &= \sum_{g \geq 0} (-s/2)^{3(1-g)} F^g(\widehat{\Sigma}) \end{aligned} \quad (3-18)$$

3.3 Existence of the $1/N$ expansion

In our case, $\mathbf{L}(X, S)$ has only one pole, at $X = \infty$, which is already present in $\mathbf{L}^{(0)}(X, S)$. We define $Y(x)$ by Eqn. 3-13 with the choice of the branch of the square root imposed by the $X \rightarrow \infty$ asymptotics in Eqn. 3-9:

$$Y(X) = -4iX \sqrt{X^2 + \frac{S}{2}}, \quad Y(x) \underset{X \rightarrow +\infty}{\sim} -4iX^2 \quad (3-19)$$

In order to identify $\mathcal{F}(\Sigma_\infty)$ computed in Eqn. 3-18 with the $s \rightarrow -\infty$ asymptotic of the τ function of the initial system (Eqn 3-6), we have to check that the assumptions announced in Sections 1.3-1.5 hold.

Remark 3.1 Σ_∞ is a regular genus 0 spectral curve: it has two simple branchpoints, at $X = \pm \sqrt{-S/2}$. It admits a rational parametrization:

$$X(z) = \gamma \left(z + \frac{1}{z} \right), \quad Y(z) = \frac{S}{2} \left(z^2 - \frac{1}{z^2} \right) \quad (3-20)$$

where $\gamma = \sqrt{\frac{-S}{8}}$. Notice that it is not the spectral curve of a matrix model, since there is no $1/z$ term in Y . This is rather the curve of the limit of a matrix model.

Lemma 3.1 $\psi(X, S)$ (resp. $\bar{\psi}(X, S)$, $\phi(X, S)$, $\bar{\phi}(X, S)$) admits a $1/N$ expansion:

$$\psi(X, S) = \left(\tilde{\psi}^{(0)}(X, S) + \sum_{l \geq 1} N^{-l} \tilde{\psi}^{(l)}(X, S) \right) \exp \left[iN \left(\frac{4}{3} X^3 + SX \right) \right] \quad (3-21)$$

and for all $l \geq 1$, $\tilde{\psi}^{(l)}(X, S)$ has only poles at $X = \pm \sqrt{\frac{-S}{2}}$ (the branchpoints of Σ_∞). In particular, it does not have singularities at $X = 0$, the other zero of Y .

proof:

We already know that $Q^{(0)}(S) = \sqrt{\frac{-S}{2}}$, and Eqn. 3-11 implies that Q has a $1/N^2$ expansion:

$$Q(S) = Q^{(0)}(S) + \sum_{l \geq 1} N^{-2l} Q^{(l)}(S) \quad (3-22)$$

Hence, \mathbf{L} and \mathbf{M} have a $1/N$ expansion. Let us apply the discussion of Section 2.3, where S plays the role of the time t . The large N limit spectral curve Σ_∞ associated to the system $\frac{1}{N} \partial_X \Psi = \mathbf{L} \Psi$ is:

$$Y(X) = \pm 4iX \sqrt{X^2 + S/2} \quad (3-23)$$

and ψ to leading order is given by:

$$\begin{aligned}\psi^{(0)}(X, S) &= \text{cte}^{(0)}(S) \sqrt{\frac{b^{(0)}(X, S)}{2Y(X)}} \exp \left[-N \int^X Y(\xi) d\xi \right] \\ &= \text{cte}^{(0)}(S) \left(\frac{-S/2}{X^2 + S/2} \right)^{1/4} \exp \left[\mp iN \left(\frac{4}{3} X^3 + SX \right) \right]\end{aligned}\quad (3-24)$$

To agree with the asymptotic of Eqn. 3-9, we must choose the minus sign. On the other hand, the large N limit spectral curve \underline{Y}_∞ associated to the system $\frac{1}{N} \partial_S \Psi = \mathbf{M} \Psi$ is:

$$\underline{Y}(X) = \pm \sqrt{-\det \mathbf{M}^{(0)}} = \pm i \sqrt{X^2 + S/2} \quad (3-25)$$

Hence, $Y(X)$ and $\underline{Y}(X)$ have no common zeroes (apart from the branchpoints $X = \pm \sqrt{-S/2}$). So, to all orders in $1/N$, $\tilde{\psi}$ cannot have singularities at $X = 0$. The same discussion would hold for ϕ , $\bar{\psi}$ and $\bar{\phi}$. \square

Corollary 3.1 $\mathcal{W}_n(X_1, \dots, X_n)$ admits a $1/N^2$ expansion:

$$\mathcal{W}_n(X_1, \dots, X_n) = \sum_{g \geq 0} N^{2-2g-n} \mathcal{W}_n^g(X_1, \dots, X_n) \quad (3-26)$$

and the singularities of $\mathcal{W}_n^g(X_1, \dots, X_n)$ away from $X_i = \infty$ are found only at branch-points $X_i = \sqrt{-S/2}$.

proof:

There exists at least an expansion in $1/N$, and the position of the singularities at all orders is a consequence of Lemma 3.1. We have seen in Thm. 2.1 that \mathcal{W}_n starts as $O(N^{2-n})$. It remains to prove that \mathcal{W}_n has parity $(-1)^n$ in N . Let us stress the dependence in N by writing \mathbf{L}_N , and Ψ_N for the solution of Eqn. 3-9. We observe that ${}^t\mathbf{L}_{-N} = \mathbf{L}_N$, which implies that ${}^t\Psi_{-N}^{-1}$ obey the same differential system as Ψ . Moreover, ${}^t\Psi_{-N}^{-1}$ has the same asymptotic behavior near $x \rightarrow \infty$ as Ψ_N , and is also of determinant 1. So, ${}^t\Psi_{-N}^{-1} = \Psi_N$, and at the level of the integrable kernel:

$$\begin{aligned}\mathcal{K}_N(x_1, x_2) &= \frac{\psi_N(x_1) \bar{\phi}(x_2) - \bar{\psi}_N(x_1) \phi(x_2)}{x_1 - x_2} \\ &= \frac{\bar{\phi}_{-N}(x_1) \psi_{-N}(x_2) - \phi_{-N}(x_1) \bar{\psi}_{-N}(x_2)}{x_1 - x_2} \\ &= \mathcal{K}_{-N}(x_2, x_1)\end{aligned}$$

But, we can see on definition (Eqn. 1-17) that the correlators take a $(-1)^n$ sign if we revert the orientation of all the cycles. Thus:

$$W_n(x_1, \dots, x_n)_{-N} = (-1)^n W_n(x_1, \dots, x_n)_N \quad (3-27)$$

□

Accordingly, we can apply Thm. 1.1, which says:

$$\begin{aligned}\mathcal{W}_n^{(g)}(X(z_1), \dots, X(z_n)) dx(z_1) \cdots dx(z_n) &= \omega_n^g(\Sigma_\infty)(z_1, \dots, z_n) \\ &= (-S/2)^{3(1-g-n/2)} \omega_n^g(\widehat{\Sigma})(z_1, \dots, z_n)\end{aligned}\tag{3-28}$$

3.4 Tau-function and symplectic invariants

The next step is the computation of the τ -function. Let us do the computation directly, as an illustration of the general proof given in Section 1.5. The τ -function associated to the unique solution of the system satisfying Eqn. 3-9 has:

$$\mathbf{T}_\infty(x, s) = iN \left(\frac{4}{3} X^3 + SX \right) \sigma_3 \tag{3-29}$$

This is correct for any N , we see on this example that \mathbf{T}_∞ is indeed given by its large N limit. The τ -function of Jimbo-Miwa-Ueno is given by (Eqn. 1-39):

$$\partial_\sigma \ln \tau = 2iN \operatorname{Res}_{X \rightarrow \infty} dX X \mathcal{W}_1(X) \tag{3-30}$$

Let us compare with the variation of $\mathcal{F}(\Sigma_\infty)$ wrt S , given by Eqn. 1-3. We have to represent $\partial_S Y dX$ with the Bergman kernel:

$$B(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2} \tag{3-31}$$

We write:

$$\begin{aligned}(\partial_S Y dX)(z) &= \frac{-i X(z) dX(z)}{\sqrt{X^2(z) + S/2}} = -i dz \left(\sqrt{X^2(z) + \frac{S}{2}} \right) \\ &= -i \operatorname{Res}_{\zeta \rightarrow z} B(z, \zeta) \sqrt{X^2(\zeta) + \frac{S}{2}} = -i \gamma \operatorname{Res}_{\zeta \rightarrow z} B(z, \zeta) \left(\zeta - \frac{1}{\zeta} \right) \\ &= i \gamma \operatorname{Res}_{\zeta \rightarrow 0, \infty} B(z, \zeta) \left(\zeta - \frac{1}{\zeta} \right)\end{aligned}$$

Thus:

$$\partial_S F^g(\Sigma_\infty) = i \gamma \operatorname{Res}_{\zeta \rightarrow 0, \infty} \left(\zeta - \frac{1}{\zeta} \right) W_1^g(X(\zeta)) dX(\zeta) \tag{3-32}$$

Σ_∞ is an hyperelliptic curve, with involution $\zeta \mapsto \frac{1}{\zeta}$. From construction in the topological recursion, we know that:

$$\mathcal{W}_1^g(X(\zeta)) + \mathcal{W}_1^g(X(1/\zeta)) = (2y)_+(X(\zeta)) \tag{3-33}$$

where y_+ is the divergent part of $Y(X)$ when $X \rightarrow +\infty$:

$$\begin{aligned} y_+ &= \left(-4iX \sqrt{X^2 + \frac{S}{2}} \right)_+ = -i(4X^2 + S) \\ &= \frac{iS}{2} \left(\zeta^2 + \frac{1}{\zeta^2} \right) \end{aligned}$$

With the last expression, one may check that adding y_+ to W_1^g in Eqn. 3-32 do not change the result. In a first step, we may replace W_1^g by:

$$\check{W}_1^g = \mathcal{W}_1^g - \delta_{g,0} y_+ \quad (3-34)$$

in Eqn. 3-32. Now, we have the symmetry $\check{W}_1^g \rightarrow -\check{W}_1^g$ when $\zeta \rightarrow 1/\zeta$. This implies that the residues at 0 and ∞ are equal:

$$\partial_S F^g(\Sigma_\infty) = 2i\gamma \operatorname{Res}_{\zeta \rightarrow \infty} \left(\zeta - \frac{1}{\zeta} \right) \check{W}_1^g(X(\zeta)) dX(\zeta) \quad (3-35)$$

In a second step, we can go back to \mathcal{W}_1^g :

$$\partial_S F^g(\Sigma_\infty) = 2i\gamma \operatorname{Res}_{\zeta \rightarrow \infty} \left(\zeta - \frac{1}{\zeta} \right) \mathcal{W}_1^g(\Sigma_\infty)(X(\zeta)) dX(\zeta) \quad (3-36)$$

Another property of \mathcal{W}_1^g from construction is that it behaves as $O(1/\zeta)$ (and even $O(1/\zeta^2)$ for $g \geq 1$) when $\zeta \rightarrow \infty$. Accordingly, we can replace $\left(\zeta - \frac{1}{\zeta} \right)$ by $\left(\zeta + \frac{1}{\zeta} \right)$. Eventually:

$$\begin{aligned} \partial_S F^g(\Sigma_\infty) &= 2i\gamma \operatorname{Res}_{\zeta \rightarrow \infty} \left(\zeta + \frac{1}{\zeta} \right) W_1^g(\Sigma_\infty)(X(\zeta)) dX(\zeta) \\ &= 2i \operatorname{Res}_{X \rightarrow \infty} dX X \mathcal{W}_1^g(\Sigma_\infty) X \end{aligned}$$

And, for the full series:

$$\partial_S \mathcal{F}(\Sigma_\infty) = 2iN \operatorname{Res}_{X \rightarrow \infty} dX X \mathcal{W}_1(X) = \partial_S \ln \tau \quad (3-37)$$

3.5 Relation with the 1-matrix model with hard edge

So far, we have proved that the Tracy-Widom GUE law $F_{\text{GUE}}(s)$, defined axiomatically from the Airy kernel (Eqn. 3-1), has the following $s \rightarrow -\infty$ expansion:

$$F_{\text{GUE}}(s) = C \exp \left(\sum_{g \geq 0} (-s/2)^{3(1-g)} F^g(\widehat{\Sigma}) \right) \quad (3-38)$$

where C is a constant, and $\widehat{\Sigma}$ is the spectral curve of equation $\widehat{y}^2 = \frac{1}{4}\widehat{x}^2(\widehat{x}^2 + 4)$. It admits the rational parametrization:

$$\begin{cases} x(z) = \frac{i}{2} \left(z + \frac{1}{z} \right) \\ y(z) = \frac{i}{2} \left(-z^2 + \frac{1}{z^2} \right) \end{cases} \quad (3-39)$$

To complete the definition of the spectral curve, let us say that the physical sheet is associated to $x = +\infty$ ($z \rightarrow -i\infty$), and that we take the Bergman kernel:

$$\widehat{B}(z_1, z_2) = \frac{dz_1 dz_2}{(z_1 - z_2)^2} \quad (3-40)$$

In [7], starting from a matrix model with quadratic potential, where all eigenvalues are restricted to be smaller than a given a , we obtained heuristically the following $s \rightarrow -\infty$ expansion:

$$F_{\text{GUE}}(s) = C_{\text{TW}} \exp \left(\sum_{g \geq 0} (-s/2)^{3(1-g)} F^g(\Sigma_{\text{TW}}) \right) \quad (3-41)$$

with a computable constant $C_{\text{TW}} = 2^{1/24} e^{\zeta'(-1)}$, and Σ_{TW} is the spectral curve of equation $y^2 = x + \frac{1}{x} - 2$. It admits the rational parametrization:

$$\begin{cases} x(\zeta) = \zeta^2 \\ y(\zeta) = \zeta - \frac{1}{\zeta} \end{cases} \quad (3-42)$$

We associate the physical sheet to $\zeta \rightarrow +\infty$, and we take as Bergman kernel:

$$B_{\text{TW}}(\zeta_1, \zeta_2) = \frac{d\zeta_1 d\zeta_2}{(\zeta_1 - \zeta_2)^2} \quad (3-43)$$

Actually, these two spectral curves are related by a symplectic transformations, under which the F^g 's are invariant, as explained in Fig. 1. Thus, we have justified Eqn. 3-41, as announced in Prop 0.1.

$$\begin{array}{ccc} (\Sigma_{\text{TW}}) \begin{cases} x(\zeta) = \zeta^2 \\ y(\zeta) = \zeta - \frac{1}{\zeta} \end{cases} & \xrightarrow{(x \leftrightarrow y)} & \begin{cases} x(\zeta) = \zeta - \frac{1}{\zeta} \\ y(\zeta) = \zeta^2 \end{cases} \\ & \searrow \text{reparametrization} & \\ & \zeta = iz & \\ & \begin{cases} x(z) = i \left(z + \frac{1}{z} \right) \\ y(z) = -z^2 \end{cases} & \xrightarrow{y \rightarrow y + 1 + \frac{x^2}{2}} & \begin{cases} x(z) = i \left(z + \frac{1}{z} \right) \\ y(z) = \frac{1}{2} \left(-z^2 + \frac{1}{z^2} \right) \end{cases} \quad (\widehat{\Sigma}) \end{array}$$

Figure 1: Symplectic equivalence of Σ_{TW} and $\widehat{\Sigma}$

3.6 Remark on correlation functions

We insist on the fact that symplectic transformations do not conserve the correlation functions. A priori:

$$\omega_n^g(\Sigma_{\text{TW}}) \neq \omega_n^g(\widehat{\Sigma}) \quad (3-44)$$

Nevertheless, one knows [19] that their difference is an exact form in each variable.

It was argued in [7] that:

$$\omega_n(\Sigma_{\text{TW}})(\zeta_1, \dots, \zeta_n) \equiv \sum_{g \geq 0} (-s/2)^{3(1-g-n/2)} \omega_n^g(\Sigma_{\text{TW}})(\zeta_1, \dots, \zeta_n) \quad (3-45)$$

is the asymptotic expansion when $s \rightarrow -\infty$ of the (rescaled) correlation function of eigenvalues $\lambda_i = 2 + sN^{-2/3}(1 + \frac{\zeta_i^2}{2})$. In other words, for $s < 0$ and any Borel subsets $J_1, \dots, J_n \subseteq]-\infty, 0[$, we have:

$$\begin{aligned} & \lim_{N \rightarrow \infty} \text{Prob} [\lambda_1 \in J_1, \dots, \lambda_n \in J_n \mid \lambda_{\max} \leq 2 + sN^{-2/3}] \\ &= \oint_{K_1} \frac{d\zeta_1}{2i\pi} \cdots \oint_{K_n} \frac{d\zeta_n}{2i\pi} \omega_n(\Sigma_{\text{TW}})(\zeta_1, \dots, \zeta_n) \end{aligned} \quad (3-46)$$

in the sense of an asymptotic series when $s \rightarrow -\infty$. K_i is the image on the curve of J_i .

A priori, this is different from the determinantal point process defined with the integrable kernel:

$$\mathcal{K}(x_1, x_2) = \frac{\psi(x_1)\bar{\phi}(x_2) - \bar{\psi}(x_1)\phi(x_2)}{x_1 - x_2} \quad (3-47)$$

of the system of Eqn. 3-6.

4 Conclusion

Tracy-Widom law

Let us summarize what we have obtained.

- Near an hard edge a approaching the natural soft edge a_{\max} , the density of eigenvalues of a random hermitian matrix is (after appropriate rescaling) locally given by

$$(\Sigma_{\text{TW}}) : y = \sqrt{x + \frac{1}{x} - 2}$$

and therefore, the partition function in this regime is asymptotically given by the symplectic invariants of Σ_{TW} :

$$\ln Z(a) \sim \sum_g (1 - a/a_{\max})^{(2-2g)\frac{3}{2}} N^{2-2g} F^g(\Sigma_{\text{TW}})$$

and the eigenvalues correlation functions of the random hermitian matrix are also asymptotically given in this regime by the symplectic invariant correlators of Σ_{TW} , i.e. by the $\omega_n^{(g)}(\Sigma_{\text{TW}})$. Let us also mention that all this holds for any problem, solvable by symplectic invariants, whose spectral curve is locally given by Σ_{TW} . This is for instance the case for the statistics of plane partitions near the boundary of the liquid region [21].

- The second step is that Σ_{TW} is symplectically equivalent to the spectral curve $\widehat{\Sigma}$:

$$(\widehat{\Sigma}) : y = \frac{1}{4}x\sqrt{x^2 + 4},$$

which shows that

$$\ln Z(a) \sim \sum_g (1 - a/a_{\max})^{(2-2g)\frac{3}{2}} N^{2-2g} F^g(\widehat{\Sigma}),$$

and since $\widehat{\Sigma}$ is the classical spectral curve of a Lax pair for the Painlevé II equation, we recover, as expected, the large s expansion of Tracy-Widom law.

- However, in the symplectic invariance of Σ_{TW} and $\widehat{\Sigma}$, only the F^g 's are invariant, the correlators are not invariant.

It is not clear to us how the symplectic transformations (in particular $x \leftrightarrow y$, which is really the non trivial one at the level of the F^g) act on an integrable system of classical spectral curve $(x, y(x))$. It would be interesting to exhibit a full (N dependent) integrable system whose classical spectral curve is directly Σ_{TW} , without having to

perform a symplectic transformation. Such a system should be related to PII, because the τ function is the same as that of Eqn. 3-6, it can be expressed in terms of a solution of PII.

For β ensembles, the generalization of this approach is an open problem. There exists a version of the topological recursion and symplectic invariants for all $\beta > 0$, computing the large N expansion in the β random matrix ensembles [10] provided that the expansion exists. Yet, the analog of Section 1.3 for $\beta \neq 2$ is missing: starting from an integrable system, it is not known how to define quantities $\mathcal{W}_n^{(\beta)}$ satisfying the loop equations of β ensembles. One could expect that β ensembles should be related to quantum integrable systems, having to do with Bethe ansatz, instead of classical Lax pair systems.

Loop equations and integrable systems

We have reviewed that it is possible to associate loop equations (or Virasoro constraints) to any first order 2×2 rational differential integrable system. In fact, we have completed the work of [3] by:

- Showing that hypothesis on analytical properties and $1/N$ expansion of the n -point functions need only to be checked for the 1-point function.
- Clarifying when these analytical properties and expansion properties holds.
- Under all these assumptions, providing a proof of the former claim that the summation of symplectic invariants F^g reconstruct the Tau function.

To fulfill this program, it was essential to include the differential system in an isomonodromic problem with a full set of times. It is in fact possible to associate loop equations to first order system of size $d \times d$, and we are currently working on the generalization of the same theory to these systems [8].

A A counterexample to the $1/N^2$ expansion

Consider a system $1/N \partial_x \Psi = \mathbf{L} \Psi$. The first loop equation is:

$$\mathcal{W}_2(x, x) + (\mathcal{W}_1(x))^2 = N^2 P_1(x) \quad (1-1)$$

where $P_1(x)$ is a rational fraction given by:

$$P_1 = -\det \mathbf{L} = \frac{1}{2} \text{Tr} \mathbf{L}^2 \quad (1-2)$$

Let us assume that $\mathbf{L} = \mathbf{L}^{(0)} + \frac{1}{N} \mathbf{L}^{(1)} + O(1/N^2)$. Then:

$$P_1 = \frac{1}{2} \text{Tr} \mathbf{L}^{(0)2} + \frac{1}{N} \text{Tr} \mathbf{L}^{(0)} \mathbf{L}^{(1)} + O(1/N^2) \quad (1-3)$$

We want to find a case where the \mathcal{W}_n do not have an expansion with fixed parity of $1/N$. It is enough to exhibit a system for which the first two terms in P_1 do not vanish.

We call Painlevé II_α the equation:

$$q''(s) = 2q(s)^3 + sq(s) - \alpha \quad (1-4)$$

where α is a fixed parameter. This equation appears as the compatibility condition of the Lax system [22]:

$$\begin{cases} \partial_x \Psi = \mathbf{L} \Psi \\ \partial_s \Psi = \mathbf{M} \Psi \end{cases}$$

$$\mathbf{L}(x, s) = \begin{pmatrix} -4ix^2 - i(s + 2q^2(s)) & 4xq(s) + 2iq'(s) + \frac{\alpha}{x} \\ 4xq(s) - 2iq'(s) + \frac{\alpha}{x} & 4ix^2 + i(s + 2q^2(s)) \end{pmatrix}$$

$$\mathbf{M}(x, s) = \begin{pmatrix} -ix & q(s) \\ q(s) & ix \end{pmatrix}$$

Let us do the rescaling:

$$\begin{cases} x = N^{1/3} X \\ s = N^{2/3} S \end{cases}, \quad q(s) = N^{1/3} Q(S) \quad (1-5)$$

This leads us to the compatibility equation:

$$\frac{1}{N^2} Q''(S) = 2Q(S)^3 + SQ(S) - \frac{\alpha}{N} \quad (1-6)$$

and the Lax system:

$$\begin{cases} \frac{1}{N} \partial_X \Psi = \mathbf{L} \Psi \\ \frac{1}{N} \partial_S \Psi = \mathbf{M} \Psi \end{cases}$$

$$\mathbf{L}(X, S) = \begin{pmatrix} -4iX^2 - i(S + Q^2(S)) & 4XQ(S) + \frac{2iQ'(S)}{N} + \frac{\alpha}{NX} \\ 4XQ(S) - \frac{2iQ'(S)}{N} + \frac{\alpha}{NX} & 4iX^2 + i(S + 2Q^2(S)) \end{pmatrix}$$

$$\mathbf{M}(X, S) = \begin{pmatrix} -iX & Q(S) \\ Q(S) & iX \end{pmatrix}$$

When $N \rightarrow \infty$, we have to the first two orders:

$$Q(S) = \sqrt{-S/2} + \frac{1}{N} \frac{\alpha}{2S} + O(1/N^2) \quad (1-7)$$

We may compute:

$$\begin{aligned} & \text{Tr}(\mathbf{L}^{(0)} \mathbf{L}^{(1)}) \\ &= \text{Tr} \begin{pmatrix} -4iX^2 & 4X\sqrt{-S/2} \\ 4X\sqrt{-S/2} & 4iX^2 \end{pmatrix} \cdot \begin{pmatrix} 0 & \frac{2X}{S} - \frac{i}{\sqrt{-2S}} + \frac{\alpha}{X} \\ \frac{2X}{S} + \frac{i}{\sqrt{-2S}} + \frac{\alpha}{X} & 0 \end{pmatrix} \\ &\neq 0 \end{aligned}$$

Hence, the fixed parity property is not satisfied. Yet, one could also make the choice to rescale α in $A = \alpha/N$, and keep A fixed. In this case, we could reproduce the argument of Section 3.3, and check the fixed parity property. This counterexample supports the idea that a $1/N^2$ expansion may exist only in a good choice of rescaling with N of the parameters of the differential system.

B Topological recursion and symplectic invariants

With the data of a spectral curve (see Section 1.1), the topological recursion defines a tower of differential forms as follows:

Definition B.1 *We define:*

$$\omega_1^{(0)}(z) = -y(z) dx(z) \quad (2-1)$$

$$\omega_2^{(0)}(z_1, z_2) = B(z_1, z_2) \quad (2-2)$$

and by recursion on $2g - 2 + n$:

$$\begin{aligned} \omega_{n+1}^{(g)}(\Sigma)(z_0, \overbrace{z_1, \dots, z_n}^J) &= \sum_i \text{Res}_{z \rightarrow a_i} K(z_0, z) \left[\omega_{n+2}^{(g-1)}(\Sigma)(z, \bar{z}, J) \right. \\ &\quad \left. + \sum_{h=0}^{g-1} \sum_{I \subseteq J}' \omega_{1+\#I}^{(h)}(\Sigma)(z, I) \omega_{1+n-\#I}^{(g-h)}(\Sigma)(\bar{z}, J \setminus I) \right] \quad (2-3) \end{aligned}$$

where the sum over i is over all branchpoints a_i (the zeroes of $dx(z)$), residues are taken as contour integrals on the Riemann surface \mathcal{C} , the kernel $K(z_0, z)$ is:

$$K(z_0, z) = -\frac{\int_{z'=\bar{z}}^z \omega_2^{(0)}(\Sigma)(z_0, z')}{2(\omega_1^{(0)}(\Sigma)(z) - \omega_1^{(0)}(\Sigma)(\bar{z}))} \quad (2-4)$$

and in the sum $\sum_h \sum_{I \subseteq J}'$ the prime $'$ means that we exclude the pairs $(h = 0, I = \emptyset)$ and $(h = g, I = J)$ from the sum.

An important property is that for $2g - 2 + n > 0$, $\omega_n^{(g)}$ is a meromorphic form in each z_i , with poles only at branchpoints, and is symmetric in all z_i 's. For instance, we have:

$$\omega_1^{(1)}(z_0) = \sum_i \operatorname{Res}_{z \rightarrow a_i} K(z_0, z) B(z, \bar{z}) \quad (2-5)$$

$$\omega_3^{(0)}(z_0, z_1, z_2) = \sum_i \operatorname{Res}_{z \rightarrow a_i} K(z_0, z) (B(z, z_1)B(\bar{z}, z_2) + B(z, z_2)B(\bar{z}, z_1)) \quad (2-6)$$

Then, having defined $\omega_n^{(g)}$ for $n \geq 1$:

Definition B.2 For $g \geq 2$, we define the symplectic invariant $\omega_0^{(g)}(\Sigma)$ also denoted $F^g(\Sigma)$ as:

$$F^g(\Sigma) = \omega_0^{(g)}(\Sigma) = \frac{1}{2 - 2g} \sum_i \operatorname{Res}_{z \rightarrow a_i} \hat{\Phi}(z) \omega_1^{(g)}(\Sigma)(z) \quad (2-7)$$

where

$$\hat{\Phi}(z) = \int_o^z y(z') dx(z') \quad (2-8)$$

and the value of F^g is independent of the base point o and path used to compute the integral.

The definitions of $F^0(\Sigma)$ and $F^1(\Sigma)$ are more involved and we refer the reader to [19].

References

- [1] O. Babelon, D. Bernard, M. Talon, *Introduction to classical integrable systems*, Cambridge Monographs on Mathematical Physics (2003)
- [2] J. Baik, R. Buckingham, J. DiFranco, *Asymptotics of Tracy-Widom distributions and the total integral of a Painlevé II function*, Comm. Math. Phys. **280** 2, pp 463-497 (2008), [math.FA/0704.3636](#)
- [3] M. Bergère, B. Eynard, *Determinantal formulae and loop equations*, [math-ph/0901.3273](#) (2009)
- [4] M. Bergère, B. Eynard, *Universal scaling limits of matrix models, and (p, q) Liouville gravity*, [math-ph/0909.0854](#) (2009)
- [5] P. Bleher, A. Its, *Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model*, Ann. Math. **150** pp 185-266 (1999)
- [6] A. Borodin, P. Deift, *Fredholm determinants, Jimbo-Miwa-Ueno tau-functions, and representation theory*, Comm. Pure App. Math. **55** 9, pp 11601230 (2002), [math-ph/0111007](#)
- [7] G. Borot, B. Eynard, S.N. Majumdar, C. Nadal, *Large deviations of the maximal eigenvalue of random matrices*, submitted to J.Stat.Phys, [math-ph/1009.1945](#) (2010)
- [8] G. Borot, B. Eynard, *Lax pairs and loop equations*, in preparation.
- [9] M. Cafasso, O. Marchal, *Double scaling limits of random matrices and minimal $(2m, 1)$ models: the merging of two cuts in a degenerate case*, [math-ph/1002.3347](#) (2010)
- [10] L. Chekhov, B. Eynard, *Matrix eigenvalue model: Feynman graph technique for all genera*, JHEP 0612:026 (2006), [math-ph/0604014](#)
- [11] D.S. Dean, S.N. Majumdar, *Large deviations of extreme eigenvalues of random matrices*, Phys. Rev. Lett. **97**, pp 160-201 (2006), [cond-mat/0609651](#)
- [12] P. Deift, A. Its, I. Krasovsky, *Asymptotics of the Airy-kernel determinant*, [math.FA/0609451](#) (2006)

- [13] P. Deift, T. Kriecherbauer, K.T-R. McLaughlin, S. Venakides, X. Zhou, *Asymptotics for polynomials orthogonal with respect to varying exponential weights*, Int. Math. Res. Notices **16**, pp 759-782 (1997)
- [14] P. Deift, T. Kriecherbauer, K.T-R. McLaughlin, S. Venakides, X. Zhou, *Strong asymptotics of orthogonal polynomials with respect to exponential weights via Riemann-Hilbert techniques*, Comm. Pure Appl. Math. **52** 12, pp 1491-1552 (1999)
- [15] P. Deift, T. Kriecherbauer, K.T-R. McLaughlin, S. Venakides, and X. Zhou. *Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory*, Comm. Pure Appl. Math. **52** 11, pp 1335-1425 (1999)
- [16] N.M. Ercolani, K.T-R. McLaughlin, *Asymptotics of the partition function for random matrices via Riemann-Hilbert techniques, and applications to graphical enumeration*, Int. Math. Res. Not. **14**, pp 755-820 (2003), [math-ph/0211022](#)
- [17] B. Eynard, *Topological expansion for the 1-hermitian matrix model correlation functions*, JHEP 024A:0904, [hep-th/0407261](#)
- [18] B. Eynard, N. Orantin, *Invariants of algebraic curves and topological recursion*, [math-ph/0702045](#) (2007)
- [19] B. Eynard, N. Orantin, *Algebraic methods in random matrices and enumerative geometry*, [math-ph/0811.3531](#) (2008)
- [20] B. Eynard, *Large N expansion of convergent matrix integrals, holomorphic anomalies, and background independence*, JHEP03(2009)003, [math-ph/0802.1788](#)
- [21] B. Eynard, *A matrix model for plane partitions*, J. Stat. Mech. 0910:P10011 (2009), [math-ph/0905.0535](#)
- [22] H. Flaschka, A.C. Newell, *Monodromy- and spectrum-preserving deformations, I*, Comm. Math. Phys. Volume **76** 1, pp 65-116 (1980), available [here](#)
- [23] A. Fokas, A. Its, A. Kapaev, V. Novokshenov, *Painlevé transcendents: the Riemann-Hilbert approach*, Mathematical Surveys and Monographs, Vol. 128 (2006)
- [24] P.J. Forrester, N.S. Witte, *Application of the τ -Function Theory of Painlevé Equations to Random Matrices: PIV, PII and the GUE*, Comm. Math. Phys. **219**, pp 357-398 (2001), [math-ph/0103025](#)

- [25] S.P. Hastings, J.B. McLeod, *A boundary value problem associated with the second Painlevé transcendent and the Korteweg-deVries equation*, Archive for Rational Mechanics and Analysis, Volume **73**, Issue 1, pp 31-51, available [here](#)
- [26] M. Jimbo, T. Miwa, K. Ueno, *Monodromy preserving deformation of linear ordinary differential equations with rational coefficients : I. General theory and τ -function*, Physica D, Volume **2**, Issue 2, pp 306-352 (1981)
- [27] D. Korotkin, H. Samtleben, *Generalization of Okamoto's equation to arbitrary 2×2 Schlesinger system*, Adv. Math. Phys. 461860 (2009) [nlin.SI/0906.1962](#)
- [28] M.L. Mehta, *Random matrices*, 3rd edition, Pure and Applied Mathematics, Vol. 142, 3rd edn (Amsterdam: Elsevier/Academic)
- [29] K. Okamoto, *On the τ -function of the Painlevé equations*, Physica D, Nonlinear Phenomena, Volume **2**, Issue 3, pp 525-535 (1981)
- [30] J.A. Ramírez, B. Rider, B. Virág, *Beta ensembles, stochastic Airy spectrum, and a diffusion*, [math-ph/0607331](#) (2006)
- [31] S. Albeverio, L. Pastur, M. Shcherbina, *On the $1/N$ expansion for some unitary invariant ensembles of random matrices*, Comm. Math. Phys. **224**, pp 271-305 (2001), available [here](#)
- [32] C. Tracy, H. Widom, *Level spacing distributions and the Airy kernel*, Comm. Math. Phys. **159** pp 151-174 (1994), [hep-th/9211141](#)